

RC  
268.5  
U55  
no. 164  
1979

Cancer Institute  
**NOGENESIS**  
Report Series

**BIOASSAY OF  
N-NITROSODIPHENYLAMINE  
FOR POSSIBLE CARCINOGENICITY**

**CAS No. 86-30-6**

**NCI-CG-TR-164**

U.S. DEPARTMENT OF HEALTH, EDUCATION, AND WELFARE  
Public Health Service  
National Institutes of Health



RC

268.5

U55

no. 164

1979

BIOASSAY OF  
N-NITROSODIPHENYLAMINE  
FOR POSSIBLE CARCINOGENICITY

Carcinogenesis Testing Program  
Division of Cancer Cause and Prevention  
National Cancer Institute  
National Institutes of Health  
Bethesda, Maryland 20205

U. S. DEPARTMENT OF HEALTH, EDUCATION, AND WELFARE  
Public Health Service  
National Institutes of Health

NIH Publication No. 79-1720



BIOASSAY OF  
N-NITROSODIPHENYLAMINE  
FOR POSSIBLE CARCINOGENICITY

Carcinogenesis Testing Program  
Division of Cancer Cause and Prevention  
National Cancer Institute  
National Institutes of Health

FOREWORD: This report presents the results of the bioassay of N-nitrosodiphenylamine conducted for the Carcinogenesis Testing Program, Division of Cancer Cause and Prevention, National Cancer Institute (NCI), National Institutes of Health, Bethesda, Maryland. This is one of a series of experiments designed to determine whether selected chemicals have the capacity to produce cancer in animals. A negative result, in which the test animals do not have a greater incidence of cancer than control animals, does not necessarily mean that a test chemical is not a carcinogen, inasmuch as the experiments are conducted under a limited set of circumstances. A positive result demonstrates that a test chemical is carcinogenic for animals under the conditions of the test and indicates that exposure to the chemical is a potential risk to man. The actual determination of the risk to man from chemicals found to be carcinogenic in animals requires a wider analysis.

CONTRIBUTORS: This bioassay of N-nitrosodiphenylamine was conducted at the NCI Frederick Cancer Research Center (FCRC) (1), Frederick, Maryland, operated for NCI (2) by Litton Bionetics, Inc.

The manager of the bioassay at FCRC was Dr. B. Ulland, the toxicologist was Dr. E. Gordon, and Drs. R. Cardy and D. Creasia compiled the data. Ms. S. Toms was responsible for management of data, Mr. D. Cameron for management of histopathology, Mr. L. Callahan for management of the computer branch, and Mr. R. Cypher for management of the facilities. Mr. A. Butler performed the computer services. Histopathologic evaluations for rats and mice were performed by Dr. P. K. Hildebrandt (3). The diagnoses included in this report represent his interpretations.

Animal pathology tables and survival tables were compiled at EG&G Mason Research Institute (4). Statistical analyses were performed by Dr. J. R. Joiner (5) and Ms. P. L. Yong (5), using methods selected for the bioassay program by Dr. J. J. Gart (6). The chemicals used in this bioassay were analyzed at FCRC by Dr. W. Zielinsky (1). The chemical narrative and analyses were reviewed and approved by Dr. W. Lijinsky (1).

This report was prepared at Tracor Jitco (5) under the direction of NCI. Those responsible for the report at Tracor Jitco were Dr. C. R. Angel, Acting Director of the Bioassay Program; Dr. S. S. Olin, Deputy Director for Science; Dr. J. F. Robens, toxicologist; Dr. R. L. Schueler, pathologist; Dr. G. L. Miller, Ms. L. A. Owen, Ms. M. S. King, and Mr. W. D. Reichardt, bioscience writers; and Dr. E. W. Gunberg, technical editor, assisted by Ms. Y. E. Presley and Ms. P. J. Graboske.

The following scientists at NCI were responsible for evaluating the bioassay experiment, interpreting the results, and reporting the findings: Dr. Kenneth C. Chu, Dr. Cipriano Cueto, Jr., Dr. J. Fielding Douglas, Dr. Richard A. Griesemer, Dr. Thomas E. Hamm, Dr. William V. Hartwell, Dr. Morton H. Levitt, Dr. Harry A. Milman, Dr. Thomas W. Orme, Dr. A. R. Patel, Dr. Sherman F. Stinson, Dr. Jerrold M. Ward, and Dr. Carrie E. Whitmire.

- 
- (1) Frederick Cancer Research Center, P.O. Box B, Frederick, Maryland.
  - (2) Carcinogenesis Testing Program, Division of Cancer Cause and Prevention, National Cancer Institute, National Institutes of Health, Bethesda, Maryland.
  - (3) Litton Bionetics, Inc., 5516 Nicholson Lane, Kensington, Maryland.
  - (4) EG&G Mason Research Institute, 1530 East Jefferson Street, Rockville, Maryland.
  - (5) Tracor Jitco, Inc., 1776 East Jefferson Street, Rockville, Maryland.
  - (6) Mathematical Statistics and Applied Mathematics Section, Biometry Branch, Field Studies and Statistics, Division of Cancer Cause and Prevention, National Cancer Institute, National Institutes of Health, Bethesda, Maryland.



## SUMMARY

A bioassay of N-nitrosodiphenylamine for possible carcinogenicity was conducted by administering the test chemical in feed to F344 rats and B6C3F1 mice.

Groups of 50 rats of each sex were administered N-nitrosodiphenylamine at one of two doses, either 1,000 or 4,000 ppm, for 100 weeks. Matched controls consisted of 20 untreated rats of each sex. All surviving rats were killed at the end of administration of the test chemical.

Groups of 50 male mice were administered N-nitrosodiphenylamine at one of two doses, either 10,000 or 20,000 ppm, for 101 weeks. Groups of 50 female mice were administered the test chemical at one of two doses, initially 5,000 or 10,000 ppm, for 38 weeks. Because of excessive depression in the amount of mean body weight gained in the dosed groups, the doses for the females were then reduced to 1,000 and 4,000 ppm, respectively, and administration at the lowered doses was continued for 60 weeks. The time-weighted average doses for the female mice were either 2,315 or 5,741 ppm. Matched controls consisted of 20 untreated mice of each sex. All surviving mice were killed at the end of administration of the test chemical.

Mean body weights of dosed rats and mice of each sex were lower than those of corresponding controls, and were dose related throughout the bioassay, except for those of female rats during the first part of the bioassay. Mortality was dose related in the female rats, but was not affected when the test chemical was administered to the male rats or the male or female mice. Survival at the end of the bioassay was 64% or greater in the dosed and control groups of rats and mice of each sex, and sufficient numbers of animals were at risk in all groups for the development of late-appearing tumors.

Transitional-cell carcinomas of the urinary bladder occurred at incidences that were dose related ( $P$  less than 0.001) in both male and female rats, and in direct comparisons the incidences of these tumors in the high-dose groups of each sex were significantly higher ( $P$  less than or equal to 0.001) than those in the corresponding controls (males: controls 0/19, low-dose 0/46, high-dose 16/45; females: controls 0/18, low-dose 0/48, high-dose 40/49). The possible mechanism by which these tumors were induced, such as calculi formation in the bladder or nitrosation of amines present in feed to a carcinogenic nitrosoamine, is unknown.

Fibromas of the integumentary system occurred in male rats at incidences that were dose related ( $P = 0.003$ ), although in direct comparisons the incidences of these tumors in the individual dosed groups were not significantly higher than those in the control group (controls 1/20, or 5%; low-dose 1/50, or 2%; high-dose 10/50, or 20%). The incidence of fibromas of the integumentary system in historical-control male F344 rats at this laboratory is 6/285, or 2%. These results suggest an association of the fibromas in the male rats with the administration of the test chemical.

No tumors occurred in the mice of either sex at incidences that were significantly higher in the dosed groups than in the corresponding control groups. The only changes related to compound administration were chronic inflammatory lesions in the urinary bladders of dosed mice.

It is concluded that under the conditions of this bioassay, N-nitrosodiphenylamine was carcinogenic for both sexes of F344 rats, inducing transitional-cell carcinomas of the urinary bladder, but was not carcinogenic for B6C3F1 mice of either sex.



## TABLE OF CONTENTS

	<u>Page</u>
I. Introduction.....	1
II. Materials and Methods.....	3
A. Chemical.....	3
B. Dietary Preparation.....	3
C. Animals.....	4
D. Animal Maintenance.....	5
E. Subchronic Studies.....	7
F. Chronic Studies.....	11
G. Clinical and Pathologic Examinations.....	11
H. Data Recording and Statistical Analyses.....	15
III. Results - Rats.....	21
A. Body Weights and Clinical Signs (Rats).....	21
B. Survival (Rats).....	21
C. Pathology (Rats).....	24
D. Statistical Analyses of Results (Rats).....	27
IV. Results - Mice.....	29
A. Body Weights and Clinical Signs (Mice).....	29
B. Survival (Mice).....	29
C. Pathology (Mice).....	32
D. Statistical Analyses of Results (Mice).....	36
V. Discussion.....	37
VI. Bibliography.....	41

## APPENDIXES

Appendix A	Summary of the Incidence of Neoplasms in Rats Administered N-Nitrosodiphenylamine in the Diet.....	45
Table A1	Summary of the Incidence of Neoplasms in Male Rats Administered N-Nitrosodiphenylamine in the Diet.....	47

		<u>Page</u>
Table A2	Summary of the Incidence of Neoplasms in Female Rats Administered N-Nitrosodiphenylamine in the Diet.....	51
Appendix B	Summary of the Incidence of Neoplasms in Mice Administered N-Nitrosodiphenylamine in the Diet .....	55
Table B1	Summary of the Incidence of Neoplasms in Male Mice Administered N-Nitrosodiphenylamine in the Diet.....	57
Table B2	Summary of the Incidence of Neoplasms in Female Mice Administered N-Nitrosodiphenylamine in the Diet.....	61
Appendix C	Summary of the Incidence of Nonneoplastic Lesions in Rats Administered N-Nitrosodiphenylamine in the Diet.....	65
Table C1	Summary of the Incidence of Nonneoplastic Lesions in Male Rats Administered N-Nitrosodiphenylamine in the Diet.....	67
Table C2	Summary of the Incidence of Nonneoplastic Lesions in Female Rats Administered N-Nitrosodiphenylamine in the Diet.....	70
Appendix D	Summary of the Incidence of Nonneoplastic Lesions in Mice Administered N-Nitrosodiphenylamine in the Diet.....	73
Table D1	Summary of the Incidence of Nonneoplastic Lesions in Male Mice Administered N-Nitrosodiphenylamine in the Diet.....	75
Table D2	Summary of the Incidence of Nonneoplastic Lesions in Female Mice Administered N-Nitrosodiphenylamine in the Diet.....	79
Appendix E	Analyses of the Incidence of Primary Tumors in Rats Administered N-Nitrosodiphenylamine in the Diet.....	83

	<u>Page</u>
Table E1      Analyses of the Incidence of Primary Tumors in Male Rats Administered N-Nitrosodiphenylamine in the Diet.....	85
Table E2      Analyses of the Incidence of Primary Tumors in Female Rats Administered N-Nitrosodiphenylamine in the Diet.....	91
Appendix F    Analyses of the Incidence of Primary Tumors in Mice Administered N-Nitrosodiphenylamine in the Diet.....	95
Table F1      Analyses of the Incidence of Primary Tumors in Male Mice Administered N-Nitrosodiphenylamine in the Diet.....	97
Table F2      Analyses of the Incidence of Primary Tumors in Female Mice Administered N-Nitrosodiphenylamine in the Diet.....	101

#### TABLES

Table 1      N-Nitrosodiphenylamine Subchronic Feeding Studies in Rats .....	8
Table 2      N-Nitrosodiphenylamine Subchronic Feeding Studies in Mice.....	9
Table 3      N-Nitrosodiphenylamine Chronic Feeding Studies in Rats.....	12
Table 4      N-Nitrosodiphenylamine Chronic Feeding Studies in Mice.....	13


#### FIGURES

Figure 1      Growth Curves for Rats Administered N-Nitrosodiphenylamine in the Diet.....	22
Figure 2      Survival Curves for Rats Administered N-Nitrosodiphenylamine in the Diet.....	23
Figure 3      Growth Curves for Mice Administered N-Nitrosodiphenylamine in the Diet.....	30

Figure 4	Survival Curves for Mice Administered N-Nitrosodiphenylamine in the Diet.....	31
----------	--	----

## I. INTRODUCTION

N-Nitrosodiphenylamine (CAS 86-30-6; NCI C02880) is a nitroso-amine which is used as a vulcanization retarder in curing natural rubber and the synthetic elastomers styrenebutadiene and nitrile-butadiene (Del Gatto, 1968; Stern, 1967). U.S. pro-



N-nitrosodiphenylamine

duction in 1976 was 1.3 million pounds (United States International Trade Commission, 1977).

The acute oral LD<sub>50</sub> for N-nitrosodiphenylamine in BD rats is estimated to be 3,000 mg/kg (Druckrey et al., 1967). The LD<sub>50</sub> of this compound in white mice (sex and strain not specified) when administered by intragastric intubation is 3,850 mg/kg (Zhilova and Kasparov, 1966).

Approximately 100 N-nitroso compounds have been demonstrated to be carcinogenic in animal systems (Magee et al., 1976) since the original work of Magee and Barnes (1956) which demonstrated the hepatocarcinogenicity of dimethylnitrosamine. Nitrosoamines have



been shown to cause cancer of the liver, lungs, esophagus, nasal cavities, bladder, and kidney in either rats or mice (Weisburger, 1975). Carcinogenic effects have also been observed in dogs, pigs, hamsters, fish, and primates (Weisburger, 1975; Magee et al., 1976).

There is strong evidence that bladder cancer rates are higher among men employed in the rubber industry than among the general population (Boyland et al., 1968; Case and Hosker, 1954), and there is a specific association between bladder cancer and the handling of chemicals by employees in shipping and receiving in the rubber industry, as well as those involved in compounding, mixing, and calendering rubber (McMichael et al., 1976).

N-nitrosodiphenylamine was tested by Innes et al. (1969) in a large-scale screen of industrial compounds for carcinogenic activity. Since the results of this preliminary bioassay in mice did not clearly associate the incidence of any tumor with administration of the test chemical, N-nitrosodiphenylamine was selected for further testing in the Carcinogenesis Testing Program.

## II. MATERIALS AND METHODS

### A. Chemical

Redax<sup>®</sup> (N-nitrosodiphenylamine) was obtained from R. T. Vanderbilt as a dark-brown solid. Its purity was estimated by high-pressure liquid chromatography to be 98%, with two impurities. Its melting point was 64.9°C, as compared with its reported melting point range of 63 to 66°C and its infrared spectrum was consistent with the chemical structure. Mass spectral analysis gave a molecular ion at 198 m/e and a base peak at 168 m/e. Elemental analysis showed 72.4% carbon, 5.2% hydrogen, and 13.6% nitrogen (theoretical 72.7%, 5.1%, and 14.1%).

### B. Dietary Preparation

Test diets containing N-nitrosodiphenylamine were prepared approximately weekly in 6- to 12-kg batches at appropriate doses. A known weight of the chemical was first mixed with an equal weight of autoclaved Wayne<sup>®</sup> Sterilizable Lab Meal with 4% fat (Allied Mills, Inc., Chicago, Ill.), using a mortar and pestle. The mixing was continued with second and third additions

of feed, and final mixing was performed with the remaining quantity of feed for a minimum of 15 minutes in a Patterson-Kelly twin-shell blender with an intensifier bar. The diets were routinely stored at 7°C until used.

### C. Animals

Male and female F344 (Fischer) rats and B6C3F1 mice were obtained as 4-week-old weanlings, all within 3 days of the same age, from the NCI Frederick Cancer Research Center animal farm (Frederick, Md.). The animals were housed within the test facility for 2 weeks and then were assigned four rats to a cage and five mice to a cage on a weight basis for each cage of animals of a given species and sex. For use in the chronic study, the male rats were required to weigh 90 to 105 g, averaging at least 100 g; the female rats, 80 to 95 g, averaging at least 90 g; the male mice, 18 to 22 g, averaging at least 19.5 g; and the female mice, 17 to 21 g, averaging at least 18.5 g. Individual animals were identified by ear punch.

#### D. Animal Maintenance

The animals were housed in polycarbonate cages (Lab Products Inc., Garfield, N.J.), 19 x 10-1/2 x 8 inches for the rats and 11-1/2 x 7-1/2 x 5 inches for the mice, which were suspended from aluminum racks (Scientific Cages, Inc., Bryan, Tex.) and were covered by nonwoven polyester-fiber 12-mil-thick filter paper (Hoeltge, Inc., Cincinnati, Ohio). The bedding used was Absorb-dri<sup>®</sup> hardwood chips (Northeastern Products, Inc., Warrenburg, N.Y.). The feed supplied was presterilized Wayne<sup>®</sup> Sterilizable Lab Meal provided ad libitum in suspended stainless steel hoppers and replenished at least three times per week. Water, acidified to pH 2.5, was supplied ad libitum from glass bottles. Sipper tubes (Lab Products, Inc.) were suspended through the tops of the cages.

The contaminated bedding was disposed of through an enclosed vacuum line that led to a holding tank from which the bedding was fed periodically into an incinerator. The cages were sanitized twice per week and the feed hoppers twice per month at 82 to 88°C in a tunnel-type cagewasher (Industrial Washing Machine Corp., Mataway, N. J.), using the detergents, Clout<sup>®</sup> (Pharmaceutical Research Laboratories, Greenwich, Conn.) or Oxford D'Chlor (Oxford Chemicals, Atlanta, Ga.).

The glass bottles and sipper tubes were sanitized at 82 to 88°C in a tunnel-type bottle washer (Consolidated Equipment Supply Co., Mercersburg, Pa.) three times per week, using a Calgen Commercial Division detergent (St. Louis, Mo.). The racks for the cages were changed and sanitized at or above 82°C in a rack washer (Consolidated Equipment Supply Co.) once per month, using the Calgen Commercial Division detergent, and the filter paper was changed at the same time.

The air in the animal rooms was maintained at 22 to 24°C and at 45 to 55% relative humidity. Fresh air was passed through a filter of 65% efficiency and a bag filter of 95% efficiency at the intake and through a "Z"-type roughing filter of 30% efficiency and a bag system of 90 to 95% efficiency at the exhaust (American Air Filters, Louisville, Ky.; Mine Safety Appliances, Pittsburgh, Pa.); the air was not recirculated. Room air was changed 15 times per hour. The air pressure was maintained negative to a clean hallway and positive to a return hallway. Fluorescent lighting was provided automatically on a 12-hour-per-day cycle.

Rats administered N-nitrosodiphenylamine and their corresponding controls were housed in the same room as rats on feeding studies of the following chemicals:



(CAS 156-62-7) calcium cyanamide  
(CAS 3165-93-3) 4-chloro-o-toluidine hydrochloride

Mice administered N-nitrosodiphenylamine and their corresponding controls were housed in the same room as mice on feeding studies of the following chemicals:

(CAS 156-62-7) calcium cyanamide  
(CAS 99-81-5) (2-chloroethyl)trimethylammonium chloride (CCC)  
(CAS 95-80-7) 2,4-diaminotoluene  
(CAS 19010-66-3) lead dimethyldithiocarbamate  
(CAS 88-96-0) phthalamide  
(CAS 120-62-7) piperonyl sulfoxide  
(CAS 137-170-7) 2,4,5-trimethylaniline

#### E. Subchronic Studies

Subchronic feeding studies were conducted to estimate the maximum tolerated doses (MTD's) of N-nitrosodiphenylamine, on the basis of which two concentrations (referred to in this report as "low" and "high" doses) were selected for administration in the chronic studies. Groups of five rats and five mice of each sex were fed diets containing N-nitrosodiphenylamine at one of several doses, while groups of five control animals of each species and sex were administered basal diet only. The length of the study in male rats was 11 weeks, while in female rats and male and female mice it was 8 weeks. Each animal was weighed twice per week. Tables 1 and 2 show the number of animals in each dosed group that survived to the end of the study, the week on study when the last

Table 1. N-Nitrosodiphenylamine Subchronic Feeding Studies in Rats

		Male		Female		
Dose (ppm)	Surviv- al (a)	Week on Study	Mean Weight	Surviv- al (a)	Week on Study	Mean Weight
		When Last Death Occurred	at Week 11 as % of Control		When Last Death Occurred	at Week 7 as % of Control
First Study						
1,000	5/5		92			
2,000	5/5		94			
3,000	5/5		92			
4,000	5/5		88			
6,000	5/5		87			
8,000	5/5		81			
10,000	5/5		84			
Second Study						
4,000				5/5		86
8,000				5/5		86
16,000				2/5	5	63
24,000				0/5	4	
32,000				0/5	4	
46,000				0/5	2	

(a) Number surviving/number in group.

Table 2. N-Nitrosodiphenylamine Subchronic Feeding Studies in Mice

Dose (ppm)	Male		Female	
	Surviv- al (a)	Mean Weight at Week 7 as % of Control	Surviv- al (a)	Mean Weight at Week 7 as % of Control
First Study				
3,160	5/5	104	5/5	95
4,640	5/5	106	5/5	88
6,800	5/5	107	5/5	96
10,000	5/5	108	5/5	97
14,700	5/5	104	5/5	93
Second Study				
4,250	5/5	97		
7,500	5/5	92		
8,500	5/5	97		
9,500	5/5	86		
11,000	5/5	90		
15,000	5/5	88		
22,000	5/5	86	5/5	94
32,000			5/5	94
46,000			5/5	86

(a) Number surviving/number in group.

death occurred, and the mean body weights of each dosed group at week 11 for the male rats and at week 7 for the female rats and the male and female mice, expressed as percentages of mean body weights of corresponding controls. Weights at week 7 rather than week 8 are included, since they were used for the MTD determination.

At the end of the subchronic studies, all animals were killed using CO<sub>2</sub> and necropsied. The only histopathologic lesions observed were trace amounts of pigmentation of Kupffer's cells in hepatic sinusoids of male mice dosed at 46,000 ppm.

A 10% depression in mean body weight was the major criterion for estimation of MTD's. The doses required to produce this response were determined by the following procedure: first, least squares regressions of mean body weights versus days on study were used to estimate mean body weights of each of the dosed groups at day 49. Next, probits of the percent weights of dosed groups at day 49 relative to weights of corresponding control groups were plotted against the logarithms of the doses, and least squares regressions fitted to the data were used to estimate the doses required to induce 10% depression in weight.

Based on the data thus obtained, the low and high doses for

chronic studies using rats were set at 1,000 ppm and 4,000 ppm; using male mice, 10,000 and 20,000 ppm, and using female mice, 5,000 and 10,000 ppm.

#### F. Chronic Studies

The test groups, doses administered, and durations of the chronic feeding studies are shown in tables 3 and 4. Due to excessive weight depression in the dosed female mice, doses for the low- and high-dose groups of the females were reduced to 1,000 and 4,000 ppm, respectively, after 38 weeks.

#### G. Clinical and Pathologic Examinations

All animals were observed twice daily, and the occurrences of sick, tumor-bearing, and moribund animals were recorded. Clinical examination and palpation for masses were performed each month, and the animals were generally weighed at least once per month, except for the period of week 42 through week 64, when weights were not recorded for the rats. Moribund animals and animals that survived to the end of the bioassay were killed using CO<sub>2</sub> and necropsied.



Table 3. N-Nitrosodiphenylamine Chronic Feeding Studies in Rats

Sex and Test Group	Initial No. of Animals(a)	N-Nitroso- diphenylamine in Diet(b) (ppm)	Time on Study (weeks)
<u>Male</u>			
Matched-Control	20	0	100
Low-Dose	50	1,000	100
High-Dose	50	4,000	100
<u>Female</u>			
Matched-Control	20	0	100
Low-Dose	50	1,000	100
High-Dose	50	4,000	100

(a) All animals were 6 weeks of age when placed on study.

(b) Test and control diets were provided ad libitum 7 days per week.

Table 4. N-Nitrosodiphenylamine Chronic Feeding Studies in Mice

<u>Sex and Test Group</u>	<u>Initial No. of Animals(a)</u>	<u>N-Nitroso- diphenylamine in Diet(b) (ppm)</u>	<u>Time on Study (weeks)</u>	<u>Time-Weighted Average Dose (c) (ppm)</u>
<u>Male</u>				
Matched-Control	20	0	101	
Low-Dose	50	10,000	101	
High-Dose	50	20,000	101	
<u>Female</u>				
Matched-Control	20	0	101	
Low-Dose(d)	50	5,000	38	2,315
		1,000	60	
High-Dose(d)	50	10,000	38	5,741
		4,000	60	

(a) All animals were 6 weeks of age when placed on study.

(b) Test and control diets were provided ad libitum 7 days per week.

(c) Time-weighted average dose =  $\frac{\Sigma(\text{dose in ppm} \times \text{no. of weeks at that dose})}{\Sigma(\text{no. of weeks receiving each dose})}$

(d) Because the dosed female mice failed to gain in weight comparable to the controls, the administration of doses to the females was stopped after 38 weeks and was started again at the lower doses indicated after 41 weeks.

The pathologic evaluation consisted of gross and microscopic examination of major tissues, major organs, and all gross lesions. The tissues were preserved in 10% buffered formalin, embedded in paraffin, sectioned, and stained with hematoxylin and eosin. The following tissues were examined microscopically: skin, lungs and bronchi, trachea, bone marrow (femur), spleen, lymph nodes (mesenteric and submandibular), thymus, heart, salivary glands (parotid, sublingual, and submaxillary), liver, pancreas, esophagus, stomach (glandular and nonglandular), small and large intestine, kidney, urinary bladder, pituitary, adrenal, thyroid, parathyroid, testis, prostate, mammary gland, uterus, ovary, brain (cerebrum and cerebellum), and all tissue masses. Peripheral blood smears also were made for all animals, whenever possible.

Necropsies were also performed on all animals found dead, unless precluded in whole or in part by autolysis or cannibalization. Thus, the number of animals from which particular organs or tissues were examined microscopically varies and does not necessarily represent the number of animals that were placed on study in each group.

## H. Data Recording and Statistical Analyses

Pertinent data on this experiment have been recorded in an automatic data processing system, the Carcinogenesis Bioassay Data System (Linhart et al., 1974). The data elements include descriptive information on the chemicals, animals, experimental design, clinical observations, survival, body weight, and individual pathologic results, as recommended by the International Union Against Cancer (Berenblum, 1969). Data tables were generated for verification of data transcription and for statistical review.

These data were analyzed using the appropriate statistical techniques described in this section. Those analyses of the experimental results that bear on the possibility of carcinogenicity are discussed in the statistical narrative sections.

Probabilities of survival were estimated by the product-limit procedure of Kaplan and Meier (1958) and are presented in this report in the form of graphs. Animals were statistically censored as of the time that they died of other than natural causes or were found to be missing; animals dying from natural causes were not statistically censored. Statistical analyses for a possible dose-related effect on survival used the method of Cox

(1972) for testing two groups for equality and Tarone's (1975) extensions of Cox methods for testing for a dose-related trend. One-tailed P values have been reported for all tests except the departure from linearity test, which is only reported when its two-tailed P value is less than 0.05.

The incidence of neoplastic or nonneoplastic lesions has been given as the ratio of the number of animals bearing such lesions at a specific anatomic site (numerator) to the number of animals in which that site is examined (denominator). In most instances, the denominators included only those animals for which that site was examined histologically. However, when macroscopic examination was required to detect lesions prior to histologic sampling (e.g., skin or mammary tumors), or when lesions could have appeared at multiple sites (e.g., lymphomas), the denominators consist of the numbers of animals necropsied.

The purpose of the statistical analyses of tumor incidence is to determine whether animals receiving the test chemical developed a significantly higher proportion of tumors than did the control animals. As a part of these analyses, the one-tailed Fisher exact test (Cox, 1970) was used to compare the tumor incidence of a control group with that of a group of dosed animals at each dose level. When results for a number of dosed groups (k) are



compared simultaneously with those for a control group, a correction to ensure an overall significance level of 0.05 may be made. The Bonferroni inequality (Miller, 1966) requires that the P value for any comparison be less than or equal to  $0.05/k$ . In cases where this correction was used, it is discussed in the narrative section. It is not, however, presented in the tables, where the Fisher exact P values are shown.

The Cochran-Armitage test for linear trend in proportions, with continuity correction (Armitage, 1971), was also used. Under the assumption of a linear trend, this test determines if the slope of the dose-response curve is different from zero at the one-tailed 0.05 level of significance. Unless otherwise noted, the direction of the significant trend is a positive dose relationship. This method also provides a two-tailed test of departure from linear trend.

A time-adjusted analysis was applied when numerous early deaths resulted from causes that were not associated with the formation of tumors. In this analysis, deaths that occurred before the first tumor was observed were excluded by basing the statistical tests on animals that survived at least 52 weeks, unless a tumor was found at the anatomic site of interest before week 52. When such an early tumor was found, comparisons were based exclusively

on animals that survived at least as long as the animal in which the first tumor was found. Once this reduced set of data was obtained, the standard procedures for analyses of the incidence of tumors (Fisher exact tests, Cochran-Armitage tests, etc.) were followed.

When appropriate, life-table methods were used to analyze the incidence of tumors. Curves of the proportions surviving without an observed tumor were computed as in Saffiotti et al. (1972). The week during which an animal died naturally or was sacrificed was entered as the time point of tumor observation. Cox's methods of comparing these curves were used for two groups; Tarone's extension to testing for linear trend was used for three groups. The statistical tests for the incidence of tumors which used life-table methods were one-tailed and, unless otherwise noted, in the direction of a positive dose relationship. Significant departures from linearity ( $P$  less than 0.05, two-tailed test) were also noted.

The approximate 95 percent confidence interval for the relative risk of each dosed group compared with its control was calculated from the exact interval on the odds ratio (Gart, 1971). The relative risk is defined as  $p_t/p_c$  where  $p_t$  is the true binomial probability of the incidence of a specific type of tumor

in a dosed group of animals and  $p_c$  is the true probability of the spontaneous incidence of the same type of tumor in a control group. The hypothesis of equality between the true proportion of a specific tumor in a dosed group and the proportion in a control group corresponds to a relative risk of unity. Values in excess of unity represent the condition of a larger proportion in the dosed group than in the control.

The lower and upper limits of the confidence interval of the relative risk have been included in the tables of statistical analyses. The interpretation of the limits is that in approximately 95% of a large number of identical experiments, the true ratio of the risk in a dosed group of animals to that in a control group would be within the interval calculated from the experiment. When the lower limit of the confidence interval is greater than one, it can be inferred that a statistically significant result ( $P$  less than 0.025 one-tailed test when the control incidence is not zero,  $P$  less than 0.050 when the control incidence is zero) has occurred. When the lower limit is less than unity, but the upper limit is a greater than unity, the lower limit indicates the absence of a significant result while the upper limit indicates that there is a theoretical possibility of the induction of tumors by the test chemical, which could not be detected under the conditions of this test.



### III. RESULTS - RATS

#### A. Body Weights and Clinical Signs (Rats)

Mean body weights of dosed male and female rats were lower than those of corresponding controls, and were dose related throughout the bioassay for the males, but only sometime following week 40 for the females (figure 1). Some fluctuation in the growth curves may be due to mortality; as the size of a group diminishes, the mean body weight may be subject to greater variation. No data were recorded for the period from weeks 38 to 68. Corneal opacity occurred at higher incidences in high-dose males (15/50) and low-dose females (16/50) than in corresponding control males (0/20) and control females (1/20) and may have been related to the administration of the test chemical.

#### B. Survival (Rats)

The Kaplan and Meier curves estimating the probabilities of survival for male and female rats administered N-nitrosodiphenylamine in the diet at the doses of this bioassay, together with those of the matched controls, are shown in figure 2. The







Figure 1. Growth Curves for Rats Administered N-Nitrosodiphenylamine in the Diet





Figure 2. Survival Curves for Rats Administered N-Nitrosodiphenylamine in the Diet

result of the Tarone test for dose-related trend in mortality is not significant in male rats. In females, the result of the Tarone test is significant ( $P = 0.024$ ).

In male rats, 43/50 (86%) of the high-dose group, 44/50 (88%) of the low-dose group, and 16/20 (80%) of the control group lived to the end of the study. In females, 35/50 (70%) of the high-dose group, 44/50 (88%) of the low-dose group, and 18/20 (90%) of the control group lived to the end of the study.

Sufficient numbers of rats of each sex were at risk for the development of late-appearing tumors.

### C. Pathology (Rats)

Histopathologic findings on neoplasms in rats are summarized in Appendix A, tables A1 and A2; findings on nonneoplastic lesions are summarized in Appendix C, tables C1 and C2.

A variety of neoplasms are represented in both dosed and control groups of rats. Most of the types of tumors represented have been encountered previously in control aging F344 rats. There was a high incidence of tumors of the urinary bladder in the

high-dose groups of each sex. The incidence of these tumors, along with bladder hyperplasia, are shown in the following tabulation:

	<u>Male</u>			<u>Female</u>		
	<u>Control</u>	<u>Low Dose</u>	<u>High Dose</u>	<u>Control</u>	<u>Low Dose</u>	<u>High Dose</u>
Number of Tissues Examined	19	46	45	18	48	49
Transitional-cell Carcinoma			16(36%)			40(82%)
Transitional-cell Carcinoma with Squamous Metaplasia			1(2%)			2(4%)
Epithelial Hyperplasia		2(4%)	6(13%)		4(8%)	7(14%)

In high-dose groups of each sex, the entire spectrum from transitional-cell hyperplasia to transitional-cell carcinoma was observed in the urinary bladder. The hyperplastic foci consisted of enlarged transitional cells and the epithelium was several (7 to 10) layers in thickness. In some of the hyperplastic foci there was a tendency for the cells in the basilar layer to compartmentalize and form circular, almost acinar-like structures. Mitotic figures were often noted in these basilar cells. As the epithelium increased in thickness, fibrous tissue strands began to appear, forming a connective tissue stroma for

the proliferating epithelium. These lesions were diagnosed as transitional-cell neoplasms. The epithelium covering these fibrous strands was several (7 to 10) layers thick and mitotic figures were quite numerous in most cases. Many of the tumors were similar to papillomas; however, the thickness and activity of the epithelium was consistent with papillary transitional-cell carcinoma. Many of the tumors had less fibrous stroma, the mass consisted of solid sheets of epithelial cells, or was occasionally arranged in cords. In three cases there was squamous metaplasia. The base of the tumor was narrow in many cases, but was also rather broad in many others. The degree of infiltration into deeper layers of the bladder wall was also variable. There appeared to be a tendency for the tumor mass to remain rather superficial until the mass was quite large in size and the tumor cells were more anaplastic and active. At this time there was infiltration into the deeper layers; however, in only one case was there invasion through the entire wall and beyond the serosa. In none of these animals was a transitional-cell metastatic focus seen in another organ.

A second type of tumor, fibroma of the subcutis and skin, was observed at a higher frequency in the male high-dose group (10/45) than in the male low-dose group (1/46), in the male controls (1/19), or in any of the female groups.



The fibromas were composed of well-differentiated, dense, well-circumscribed areas of fibrous tissue.

A variety of nonneoplastic lesions were represented among both control and dosed groups of animals. Such lesions have been encountered previously in untreated aging F344 rats and are not considered to be compound related.

Based on the histopathologic examination, the high incidence of bladder tumors in both male and female high-dose groups indicates that N-nitrosodiphenylamine was carcinogenic for F344 rats under the conditions of this bioassay.

#### D. Statistical Analyses of Results (Rats)

Tables E1 and E2 in Appendix E contain the statistical analyses of the incidences of those primary tumors that occurred in at least two animals of one group and at an incidence of at least 5% in one or more than one group.

In both sexes of rats, transitional-cell carcinomas of the urinary bladder occurred exclusively in the high-dose group. The result of the Cochran-Armitage test indicates a significant

positive trend ( $P$  less than 0.001) in each sex. An indicated departure from linear trend ( $P = 0.042$ ) is observed in the females, due to the steep increase in the incidence of these tumors in the high-dose group. The Fisher exact test shows that the incidence in the high-dose group is significantly higher ( $P$  less than or equal to 0.001) than that in the control group in each sex. The statistical conclusion is that the incidence of transitional-cell carcinoma of the urinary bladder in rats is related to the administration of N-nitrosodiphenylamine.

In male rats, the result of the Cochran-Armitage test for positive dose-related trend in the incidence of fibroma of the integumentary system is significant ( $P = 0.003$ ), but the results of the Fisher exact test are not significant. The historical records at this laboratory show an incidence of these tumors of 6/285 (2%), compared with 10/50 (20%) in the high-dose male rats of this bioassay.

Significant results in the negative direction are observed in the incidences of pituitary and adrenal tumors in male rats and in the incidence of hematopoietic tumors in female rats. In female rats, the significance in the negative direction may be accounted for by the difference in survival among the dosed and control groups.

#### IV. RESULTS - MICE



##### A. Body Weights and Clinical Signs (Mice)

Mean body weights of dosed male and female mice were lower than those of corresponding controls, especially for the females, and were dose related throughout the bioassay (figure 3). Some fluctuation in the growth curves may be due to mortality; as the size of a group diminishes, the mean body weight may be subject to greater variation. Tissue masses occurred at low incidences in both control and dosed groups of mice.

##### B. Survival (Mice)

The Kaplan and Meier curves estimating the probabilities of survival for male and female mice administered N-nitrosodiphenylamine in the diet at the doses of this bioassay, together with those of the matched controls, are shown in figure 4. The result of the Tarone test for dose-related trend in mortality is not significant in either sex.

In male mice, 41/50 (82%) of the high-dose group, 46/50 (92%) of



**Figure 3. Growth Curves for Mice Administered N-Nitrosodiphenylamine in the Diet**




Figure 4. Survival Curves for Mice Administered N-Nitrosodiphenylamine in the Diet



the low-dose group, and 18/20 (90%) of the control group lived to the end of the study. In females, 31/50 (62%) of the high-dose group, 42/50 (84%) of the low-dose group, and 16/20 (80%) of the control group survived to the end of the study.

Sufficient numbers of mice of each sex were at risk for the development of late-appearing tumors.

#### C. Pathology (Mice)

Histopathologic findings on neoplasms in mice are summarized in Appendix B, tables B1 and B2; findings on nonneoplastic lesions are summarized in Appendix D, tables D1 and D2.

There were similar incidences and types of tumors in control and dosed mice, and none appeared to be related to administration of the test chemical. However, there was a high incidence of bladder lesions in the dosed mice, as shown in the following tabulation:

	Male			Female		
	<u>Control</u>	<u>Low Dose</u>	<u>High Dose</u>	<u>Control</u>	<u>Low Dose</u>	<u>High Dose</u>
Number of Tissues Examined	18	49	46	18	47	38
Transitional-cell Carcinoma		1(2%)			1(2%)	
Transitional-cell Papilloma or Papilloma, NOS		1(2%)	1(2%)			
Hemangioma					1(2%)	
Epithelial Hyperplasia		2(4%)	7(15%)		3(6%)	6(16%)
Inflammation, Chronic Submucosal		12(24%)	31(67%)		31(66%)	30(79%)

The lesion most frequently observed in the bladders of the mice was chronic submucosal inflammation. No such lesion was observed in any control animal. An occasional focus of lymphocytes and an occasional blood vessel cuffed with lymphocytes, which are not an uncommon finding in the submucosa of the urinary bladder of normal mice, were observed in the control animals in this study. In the dosed animals diagnosed as having chronic submucosal inflammation of the bladder, there was an increase in the number of lymphocytes which was manifested by increased size and number of lymphocytic foci, infiltration of lymphocytes between collagen fibers, and more numerous blood vessels cuffed with lymphocytes. These lymphocytic cuffs were also much greater in thickness.

A degeneration of the collagen fibers of the submucosa was observed only in dosed mice. The degeneration was characterized by shrinking and curling of collagen bundles, and they had a more hyalinized appearance. This change appeared to occur first near, or immediately beneath, the basement membrane of the epithelium and extended to varying depths in the submucosa. Deeper in the submucosa the collagen bundles were more plump, but also more hyalinized. Blood vessels themselves also had a change which was more evident in small arterioles. This change was a thickening of the media with hyalinization of the muscle fibers. In the majority of cases this was not a severe change, but it was indeed observable. In two cases (high-dose females) this change was accompanied by acute and chronic inflammatory foci in the vessel wall and in still another case, again a high-dose female, there was fibrinoid necrosis of the vessel wall. Overall, the submucosa of the bladder seemed somewhat thickened and in two cases was considered to be edematous.

Although a few changes were observed in the epithelium of the bladder, it is somewhat suprising that more were not seen in view of the changes in the submucosa. The hyperplasia of the epithelium usually occurred as focal areas; however, in one case it occurred as diffuse hyperplasia. Two transitional-cell

carcinomas of the bladder were encountered, one of which occurred in a low-dose male, the other in a low-dose female. One transitional-cell papilloma also was seen, in a high-dose male.

A slight perivascular lymphocytic cuffing in the kidney is a normal finding in B6C3F1 mice. There were a few animals, both control and dosed, in which the degree of cuffing was considered to be greater than usual; these were diagnosed as having chronic inflammation. There was no correlation between these kidney lesions and changes in the urinary bladder.

In addition to the bladder lesions, a large number of degenerative, proliferative, and inflammatory changes were also encountered in animals of the dosed and control groups. Again no correlation could be made between incidences of lesions and administration of the test chemical.

Because the incidence of bladder neoplasms encountered in the dosed mice of this study was very low, it does not appear that N-nitrosodiphenylamine was carcinogenic for B6C3F1 mice under the conditions of this bioassay. The compound does, however, produce a nonsuppurative inflammatory response associated with a connective tissue degeneration in the submucosa of the urinary bladder in B6C3F1 mice.

#### D. Statistical Analyses of Results (Mice)

Tables F1 and F2 in Appendix F contain the statistical analyses of the incidences of those primary tumors that occurred in at least two animals of one group and at an incidence of at least 5% in one or more than one group.

The result of the Cochran-Armitage test for dose-related trend in the incidence of tumors, and the results of the Fisher exact test comparing the incidences of tumors in the dosed groups with that in the control group are not significant.

In each of the 95% confidence intervals for relative risk, shown in the tables, the value of one is included; this indicates the absence of significant positive results. It should also be noted that each of the intervals has an upper limit greater than one, indicating the theoretical possibility of the induction of tumors by N-nitrosodiphenylamine, which could not be detected under the conditions of this test.



## V. DISCUSSION

Mean body weights of dosed rats and mice of each sex were lower than those of corresponding controls, and were dose related throughout the bioassay, except for female rats during the first part of the bioassay. Mortality was dose related in the female rats, but was not affected when the test chemical was administered to the male rats or the male or female mice. Survival at the end of the bioassay was 64% or greater in all dosed and control groups of rats and mice of each sex, and sufficient numbers of animals were at risk in all groups for the development of late-appearing tumors.

In both male and female rats, transitional-cell carcinomas of the urinary bladder occurred at incidences that were dose related ( $P$  less than 0.001), and in direct comparisons the incidences of these tumors in the high-dose groups of each sex were significantly higher ( $P$  less than or equal to 0.001) than those in the corresponding controls (males: controls 0/19, low-dose 0/46, high-dose 16/45; females: controls 0/18, low-dose 0/48, high-dose 40/49). Epithelial hyperplasia of the urinary bladder occurred in both the high- and low-dose groups of each sex, and squamous metaplasia of the bladder occurred in the high-dose

groups; neither of these lesions occurred in the corresponding control groups.

Fibromas of the integumentary system occurred in male rats at incidences that were dose related ( $P = 0.003$ ), although in direct comparisons the incidences of these tumors in the individual dosed groups were not significantly higher than those in the control group (controls 1/20, or 5%; low-dose 1/50, or 2%; high-dose 10/50, or 20%). The incidence of fibromas of the integumentary system in historical-control male F344 rats at this laboratory is 6/285, or 2%. These results suggest an association of the fibromas in male rats with the administration of the test chemical.

No tumors occurred in the mice of either sex at incidences that were significantly higher in the dosed groups than in the corresponding control groups. However, submucosal inflammation of the urinary bladder occurred at high incidences in the dosed groups of mice of each sex, and epithelial hyperplasia of the bladder occurred at low incidences; neither lesion occurred in the corresponding control groups.

Previous reports on tests of the possible carcinogenicity of N-nitrosodiphenylamine have indicated that the compound was not

carcinogenic under the conditions tested. When 25 male Wistar rats were administered N-nitrosodiphenylamine by stomach tube 5 days per week for 45 weeks at doses of 1.07 mg suspended in 1 ml of 1% aqueous methylcellulose and the animals were killed at week 53, no tumors were observed on histopathologic examination of the liver, spleen, kidneys, lung, or any organs having macroscopic changes (Argus and Hoch-Ligeti, 1961). Neither were tumors observed when 20 BD rats were administered the test chemical in the diet 7 days per week for 100 weeks at doses of 120 mg/kg, or when 24 male CB rats were administered the test chemical by intraperitoneal injection in polyethylene glycol 400 solution once per week for 6 months at doses of 2.5 mg/wk and the tests were terminated after 2 years (Boyland et al., 1968). When 18 male and 18 female mice of each of two hybrids (C57BL/6 x C3H/Anf and C57BL/6 x AKR) were administered the test chemical by stomach tube daily for 3 weeks at 1,000 mg/kg body weight, then in the diet at 3,769 ppm for 18 months, no significant incidences of tumors were observed; however, reticulum-cell sarcomas were observed ( $P = 0.05$ ) when the chemical was administered by subcutaneous injection (NTIS, 1968; Innes et al., 1969). The occurrence of statistically significant incidences of transitional-cell carcinomas of the urinary bladder in the rats of the present bioassay in contrast to the apparent lack of carcinogenicity in previous studies using oral administration may

have been due to the difference in route, dose, duration of administration of the test chemical, or in the total period of observation of the dosed animals. It must be recognized that the actual mechanism by which bladder tumors were induced, such as calculi formation or nitrosation of amines present in feed to a carcinogenic nitrosamine, is unknown.

It is concluded that under the conditions of this bioassay, N-nitrosodiphenylamine was carcinogenic for both sexes of F344 rats, inducing transitional-cell carcinomas of the urinary bladder, but was not carcinogenic for B6C3F1 mice of either sex.

## VI. BIBLIOGRAPHY

Argus, M. F. and Hoch-Ligeti, C., Comparative study of the carcinogenic activity of nitrosamines. J. Natl Cancer Inst. 27:695-701, 1961.

Armitage, P., Statistical Methods in Medical Research, John Wiley & Sons, Inc., New York, 1971, pp. 362-365.

Berenblum, I., ed., Carcinogenicity Testing: A Report of the Panel on Carcinogenicity of the Cancer Research Commission of the UICC, Vol. 2. International Union Against Cancer, Geneva, 1969.

Boyland, E., Carter, R. L., Gorrod, J. W., and Roe, F. J. C., Carcinogenic properties of certain rubber additives. Europ. J. Cancer 4:233-239, 1968.

Case, R. A. M. and Hosker, M. E., Tumour of the urinary bladder as an occupational disease in the rubber industry in England and Wales. Brit. J. prev. soc. Med. 8:39, 1954.

Cox, D. R., Regression models and life tables. J. R. Statist. Soc. B:187-220, 1972.

Cox, D. R., Analysis of Binary Data, Methuen and Co., Ltd., London, 1970, pp. 48-52.

Del Gatto, J. V., Vulcanization and curing materials. In: Materials and Compounding Ingredients for Rubber, Bill Publications, Inc., New York, 1968, p. 77.

Druckrey, H., Preussmann, R., Ivankovic, S., and Schmahl, D., Organotropic carcinogenic effects of 65 different N-nitroso-compounds on BD-rats. Z. Krebsforsch. 69(103):194-195, 1967.

Druckrey, H., Preussmann, R., Schmahl, D., and Muller, M., Chemische Konstitution und carcinogene Wirkung bei Nitrosaminen. Naturwissenschaften 48:134, 1961.

Gart, J. J., The comparison of proportions: a review of significance tests, confidence limits and adjustments for stratification. Rev. Int. Statist. Inst. 39:148-169, 1971.

Innes, J. R. M., Ulland, B. M., Valerio, M. G., Petrucelli, L., Fishbein, L., Hart, E. R., Pallotta, A. J., Bates, R. R., Falk,



H. L., Gart, J. J., Klein, M., Mitchell, I., and Peters, J., Bioassay of pesticides and industrial chemicals for tumorigenicity in mice: a preliminary note. J. Natl Cancer Inst. 42(6):1101-1106, 1969.

Kaplan, E. L. and Meier, P., Nonparametric estimation from incomplete observations. J. Amer. Statist. Assoc. 53:457-481, 1958.

Linhart, M. S., Cooper, J. A., Martin, R. L., Page, N. P., and Peters, J. A. Carcinogenesis bioassay data system. J. Comp. Biomed. Res. 7:230-248, 1974.

Magee, P. N., and Barnes, J. M., The production of malignant primary hepatic tumours in the rat by feeding dimethylnitrosamine. Brit. J. Cancer 10:114-122, 1956.

Magee, P. N., Montesano, R., and Preussman, R., N-Nitroso compounds and related carcinogens. In: Chemical Carcinogens, Searle, C. E., ed., American Chemical Society, Washington, D. C., 1976, pp. 491-577.

McMichael, A. J., Andjelkovic, D. A., and Tyroler, H. A., Cancer mortality among rubber workers: an epidemiologic study. Ann. N.Y. Acad. Sci. 271:125-137, 1976.

Miller, R. G., Jr., Simultaneous Statistical Inference, McGraw-Hill Book Co., New York, 1966, pp. 6-10.

National Technical Information Service (NTIS), Evaluation of Carcinogenic, Teratogenic, and Mutagenic Activities of Selected Pesticides and Industrial Chemicals, Vol. I. Carcinogenic Study, U. S. Department of Commerce, Washington, D. C., 1968.

Saffiotti, U., Montesano, R., Sellakumar, A. R., Cefis, F., and Kaufman, D. G., Respiratory tract carcinogenesis in hamsters induced by different numbers of administrations of benzo (a) pyrene and ferric oxide. Cancer Res 32:1073-1081, 1972.

Stern, H. J., Methods of vulcanization. Antioxidants. In: Rubber: Natural and Synthetic, Palmerton Publishing Co., Inc., New York, 1967, pp. 227-236.

Tarone, R. E., Tests for trend in life table analysis. Biometrika 62 (3):679-682, 1975.

United States International Trade Commission, Rubber-processing chemicals. In: Synthetic Organic Chemicals - United States Production and Sales, 1976, USITC Publication 833, United States International Trade Commission, Washington, D. C., 1977, p. 193.

Weisburger, J. H., Chemical carcinogenesis. In: Toxicology - The Basic Science of Poisons, Casarett, L. J., and Doull, J., eds., Macmillan Publishing Co., Inc., New York, 1975, pp. 333 and 343-347.

Zhilova, N. A. and Kasparov, A. A., Comparative toxicological characteristics of anti-scorching compounds: phthalic anhydride and N-nitrosodiphenylamine (Vulkalent A). GTPZAB 10(4):60-62, 1967.

11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65  
66  
67  
68  
69  
70  
71  
72  
73  
74  
75  
76  
77  
78  
79  
80  
81  
82  
83  
84  
85  
86  
87  
88  
89  
90  
91  
92  
93  
94  
95  
96  
97  
98  
99  
100

APPENDIX A

SUMMARY OF THE INCIDENCE OF NEOPLASMS IN RATS  
ADMINISTERED N-NITROSODIPHENYLAMINE IN THE DIET





TABLE A1.

**SUMMARY OF THE INCIDENCE OF NEOPLASMS IN MALE RATS  
ADMINISTERED N-NITROSODIPHENYLAMINE IN THE DIET**

	MATCHED CONTROL	LOW DOSE	HIGH DOSE
ANIMALS INITIALLY IN STUDY	20	50	50
ANIMALS NECROPSIED	20	50	50
ANIMALS EXAMINED HISTOPATHOLOGICALLY	20	50	50
INTEGUMENTARY SYSTEM			
*SKIN	(20)	(50)	(50)
BASAL-CELL TUMOR	1 (5%)		
SEBACEOUS ADENOMA			1 (2%)
FIBROMA			1 (2%)
*SUBCUT TISSUE	(20)	(50)	(50)
BASAL-CELL TUMOR		5 (10%)	3 (6%)
SWEAT GLAND CARCINOMA			1 (2%)
FIBROMA	1 (5%)	1 (2%)	9 (18%)
HEMANGIOMA			1 (2%)
OSTEOMA		1 (2%)	
RESPIRATORY SYSTEM			
#LUNG	(20)	(49)	(49)
ALVEOLAR/BRONCHIOLAR ADENOMA			1 (2%)
OSTEOSARCOMA, INVASIVE		1 (2%)	
HEMATOPOIETIC SYSTEM			
*MULTIPLE ORGANS	(20)	(50)	(50)
ERYTHROCYTIC LEUKEMIA		1 (2%)	
GRANULOCYTIC LEUKEMIA	2 (10%)	1 (2%)	2 (4%)
#SPLEEN	(19)	(50)	(48)
HEMANGIOSARCOMA			1 (2%)
GRANULOCYTIC LEUKEMIA	1 (5%)		
#THYMUS	(14)	(46)	
OSTEOSARCOMA, INVASIVE		1 (2%)	
# NUMBER OF ANIMALS WITH TISSUE EXAMINED MICROSCOPICALLY			
* NUMBER OF ANIMALS NECROPSIED			

TABLE A1. MALE RATS: NEOPLASMS (CONTINUED)

	MATCHED CONTROL	LOW DOSE	HIGH DOSE
CIRCULATORY SYSTEM			
#HEART	(20)	(50)	(49)
SARCOMA, NOS			1 (2%)
DIGESTIVE SYSTEM			
#LIVER	(20)	(50)	(50)
HEPATOCELLULAR ADENOMA			1 (2%)
HEMANGIOSARCOMA, METASTATIC			1 (2%)
#PANCREAS	(18)	(45)	(36)
ACINAR-CELL ADENOMA	1 (6%)	1 (2%)	
URINARY SYSTEM			
#URINARY BLADDER	(19)	(46)	(45)
TRANSITIONAL-CELL CARCINOMA			16 (36%)
ENDOCRINE SYSTEM			
#PITUITARY	(18)	(47)	(47)
CHPOMOPHOBE ADENOMA	9 (50%)	9 (19%)	7 (15%)
ACIDOPHIL ADENOMA			1 (2%)
#ADRENAL	(19)	(46)	(49)
PHEOCHROMOCYTOMA	3 (16%)	6 (13%)	1 (2%)
#THYROID	(19)	(49)	(48)
FOLLICULAR-CELL ADENOMA		2 (4%)	
FOLLICULAR-CELL CARCINOMA		1 (2%)	
C-CELL ADENOMA	3 (16%)	1 (2%)	2 (4%)
#PARATHYROID	(17)	(40)	(38)
ADENOMA, NOS			1 (3%)
#PANCREATIC ISLETS	(18)	(45)	(36)
ISLET-CELL ADENOMA	2 (11%)	1 (2%)	2 (6%)
REPRODUCTIVE SYSTEM			
*MAMMARY GLAND	(20)	(50)	(50)
ADENOMA, NOS		1 (2%)	

\* NUMBER OF ANIMALS WITH TISSUE EXAMINED MICROSCOPICALLY

\* NUMBER OF ANIMALS NECROPSIED

TABLE A1. MALE RATS: NEOPLASMS (CONTINUED)

	MATCHED CONTROL	LOW DOSE	HIGH DOSE
#TESTIS INTERSTITIAL-CELL TUMOR	(19) 15 (79%)	(50) 38 (76%)	(49) 38 (78%)
NERVOUS SYSTEM			
#BRAIN ASTROCYTOMA	(19)	(47) 1 (2%)	(46)
#CEREBELLUM ASTROCYTOMA	(19)	(47) 1 (2%)	(46)
SPECIAL SENSE ORGANS			
*EYE SQUAMOUS CELL CARCINOMA	(20)	(50)	(50) 1 (2%)
MUSCULOSKELETAL SYSTEM			
NONE			
BODY CAVITIES			
NONE			
ALL OTHER SYSTEMS			
THORAX OSTEOSARCOMA		1	
ANIMAL DISPOSITION SUMMARY			
ANIMALS INITIALLY IN STUDY	20	50	50
NATURAL DEATH <sup>a</sup>	4	2	7
MORIBUND SACRIFICE		4	
SCHEDULED SACRIFICE			
ACCIDENTALLY KILLED			
TERMINAL SACRIFICE	16	44	43
ANIMAL MISSING			

<sup>a</sup> INCLUDES AUTOLYZED ANIMALS

# NUMBER OF ANIMALS WITH TISSUE EXAMINED MICROSCOPICALLY

\* NUMBER OF ANIMALS NECROPSIED

TABLE A1. MALE RATS: NEOPLASMS (CONTINUED)

	MATCHED CONTROL	LOW DOSE	HIGH DOSE
TUMOR SUMMARY			
TOTAL ANIMALS WITH PRIMARY TUMORS*	19	48	47
TOTAL PRIMARY TUMORS	38	72	91
TOTAL ANIMALS WITH BENIGN TUMORS	19	47	45
TOTAL BENIGN TUMORS	35	66	69
TOTAL ANIMALS WITH MALIGNANT TUMORS	3	5	21
TOTAL MALIGNANT TUMORS	3	6	22
TOTAL ANIMALS WITH SECONDARY TUMORS#		1	1
TOTAL SECONDARY TUMORS		2	1
TOTAL ANIMALS WITH TUMORS UNCERTAIN- BENIGN OR MALIGNANT			
TOTAL UNCERTAIN TUMORS			
TOTAL ANIMALS WITH TUMORS UNCERTAIN- PRIMARY OR METASTATIC			
TOTAL UNCERTAIN TUMORS			
* PRIMARY TUMORS: ALL TUMORS EXCEPT SECONDARY TUMORS			
# SECONDARY TUMORS: METASTATIC TUMORS OR TUMORS INVASIVE INTO AN ADJACENT ORGAN			

TABLE A2.

**SUMMARY OF THE INCIDENCE OF NEOPLASMS IN FEMALE RATS  
ADMINISTERED N-NITROSODIPHENYLAMINE IN THE DIET**

	MATCHED CONTROL	LOW DOSE	HIGH DOSE
ANIMALS INITIALLY IN STUDY	20	50	50
ANIMALS MISSING			1
ANIMALS NECROPSIED	20	50	49
ANIMALS EXAMINED HISTOPATHOLOGICALLY	20	50	49
INTEGUMENTARY SYSTEM			
*SUBCUT TISSUE	(20)	(50)	(49)
EPITHELIAL TUMOR, NOS, BENIGN			1 (2%)
BASAL-CELL TUMOR	1 (5%)	1 (2%)	3 (6%)
FIBROMA		1 (2%)	3 (6%)
FIBROSARCOMA			1 (2%)
RESPIRATORY SYSTEM			
#LUNG	(19)	(50)	(46)
ALVEOLAR/BRONCHIOLAR CARCINOMA			1 (2%)
C-CELL CARCINOMA, METASTATIC			1 (2%)
HEMATOPOIETIC SYSTEM			
*MULTIPLE ORGANS	(20)	(50)	(49)
GRANULOCYTIC LEUKEMIA	2 (10%)		
#SPLEEN	(20)	(48)	(49)
MESENCHYMOMA, BENIGN	1 (5%)		
GRANULOCYTIC LEUKEMIA	1 (5%)		
CIRCULATORY SYSTEM			
NONE			
DIGESTIVE SYSTEM			
NONE			
# NUMBER OF ANIMALS WITH TISSUE EXAMINED MICROSCOPICALLY			
* NUMBER OF ANIMALS NECROPSIED			



TABLE A2. FEMALE RATS: NEOPLASMS (CONTINUED)

	MATCHED CONTROL	LOW DOSE	HIGH DOSE
URINARY SYSTEM			
#URINARY BLADDER TRANSITIONAL-CELL CARCINOMA	(18)	(48)	(49) 40 (82%)
ENDOCRINE SYSTEM			
#PITUITARY CHROMOPHOBE ADENOMA	(19) 8 (42%)	(49) 13 (27%)	(48) 12 (25%)
#ADRENAL PHEOCHROMOCYTOMA	(19)	(50) 1 (2%)	(48)
#THYROID FOLLICULAR-CELL ADENOMA	(20)	(49) 1 (2%)	(48) 1 (2%)
FOLLICULAR-CELL CARCINOMA	1 (5%)		
C-CELL ADENOMA	1 (5%)	5 (10%)	2 (4%)
C-CELL CARCINOMA			1 (2%)
REPRODUCTIVE SYSTEM			
*MAMMARY GLAND BASAL-CELL TUMOR	(20) 1 (5%)	(50)	(49)
ADENOMA, NOS		1 (2%)	
FIBROADENOMA	4 (20%)	6 (12%)	6 (12%)
#UTERUS HEMANGIOMA	(20) 1 (5%)	(50)	(46)
#CERVIX UTERI FIBROMA	(20)	(50) 1 (2%)	(46)
NERVOUS SYSTEM			
#CEREBELLUM MENINGIOMA	(19)	(50)	(49) 1 (2%)
SPECIAL SENSE ORGANS			
NONE			
MUSCULOSKELETAL SYSTEM			
NONE			
# NUMBER OF ANIMALS WITH TISSUE EXAMINED MICROSCOPICALLY			
* NUMBER OF ANIMALS NECROPSIED			

TABLE A2. FEMALE RATS: NEOPLASMS (CONTINUED)

	MATCHED CONTROL	LOW DOSE	HIGH DOSE
BODY CAVITIES			
*ABDOMINAL CAVITY SARCOMA, NOS	(20)	(50) 1 (2%)	(49)
ALL OTHER SYSTEMS			
NONE			
ANIMAL DISPOSITION SUMMARY			
ANIMALS INITIALLY IN STUDY	20	50	50
NATURAL DEATH <sup>a</sup>		4	10
MORIBUND SACRIFICE	2	2	4
SCHEDULED SACRIFICE			
ACCIDENTALLY KILLED			
TERMINAL SACRIFICE	18	44	35
ANIMAL MISSING			1
<sup>a</sup> INCLUDES AUTOLYZED ANIMALS			
TUMOR SUMMARY			
TOTAL ANIMALS WITH PRIMARY TUMORS*	13	23	43
TOTAL PRIMARY TUMORS	21	31	72
TOTAL ANIMALS WITH BENIGN TUMORS	13	22	22
TOTAL BENIGN TUMORS	17	30	29
TOTAL ANIMALS WITH MALIGNANT TUMORS	4	1	40
TOTAL MALIGNANT TUMORS	4	1	43
TOTAL ANIMALS WITH SECONDARY TUMORS#			1
TOTAL SECONDARY TUMORS			1
TOTAL ANIMALS WITH TUMORS UNCERTAIN- BENIGN OR MALIGNANT			
TOTAL UNCERTAIN TUMORS			
TOTAL ANIMALS WITH TUMORS UNCERTAIN- PRIMARY OR METASTATIC			
TOTAL UNCERTAIN TUMORS			
* PRIMARY TUMORS: ALL TUMORS EXCEPT SECONDARY TUMORS			
# SECONDARY TUMORS: METASTATIC TUMORS OR TUMORS INVASIVE INTO AN ADJACENT ORGAN			

10  
9  
8  
7  
6  
5  
4  
3  
2  
1

## APPENDIX B

SUMMARY OF THE INCIDENCE OF NEOPLASMS IN MICE  
ADMINISTERED N-NITROSODIPHENYLAMINE IN THE DIET

11 12 13 14 15 16 17 18 19 20 21 22 23 24 25 26 27 28 29 30 31 32 33 34 35 36 37 38 39 40 41 42 43 44 45 46 47 48 49 50 51 52 53 54 55 56 57 58 59 60 61 62 63 64 65 66 67 68 69 70 71 72 73 74 75 76 77 78 79 80 81 82 83 84 85 86 87 88 89 90 91 92 93 94 95 96 97 98 99 100 101 102 103 104 105 106 107 108 109 110 111 112 113 114 115 116 117 118 119 120 121 122 123 124 125 126 127 128 129 130 131 132 133 134 135 136 137 138 139 140 141 142 143 144 145 146 147 148 149 150 151 152 153 154 155 156 157 158 159 160 161 162 163 164 165 166 167 168 169 170 171 172 173 174 175 176 177 178 179 180 181 182 183 184 185 186 187 188 189 190 191 192 193 194 195 196 197 198 199 200 201 202 203 204 205 206 207 208 209 210 211 212 213 214 215 216 217 218 219 220 221 222 223 224 225 226 227 228 229 230 231 232 233 234 235 236 237 238 239 240 241 242 243 244 245 246 247 248 249 250 251 252 253 254 255 256 257 258 259 260 261 262 263 264 265 266 267 268 269 270 271 272 273 274 275 276 277 278 279 280 281 282 283 284 285 286 287 288 289 290 291 292 293 294 295 296 297 298 299 300 301 302 303 304 305 306 307 308 309 310 311 312 313 314 315 316 317 318 319 320 321 322 323 324 325 326 327 328 329 330 331 332 333 334 335 336 337 338 339 340 341 342 343 344 345 346 347 348 349 350 351 352 353 354 355 356 357 358 359 360 361 362 363 364 365 366 367 368 369 370 371 372 373 374 375 376 377 378 379 380 381 382 383 384 385 386 387 388 389 390 391 392 393 394 395 396 397 398 399 400 401 402 403 404 405 406 407 408 409 410 411 412 413 414 415 416 417 418 419 420 421 422 423 424 425 426 427 428 429 430 431 432 433 434 435 436 437 438 439 440 441 442 443 444 445 446 447 448 449 450 451 452 453 454 455 456 457 458 459 460 461 462 463 464 465 466 467 468 469 470 471 472 473 474 475 476 477 478 479 480 481 482 483 484 485 486 487 488 489 490 491 492 493 494 495 496 497 498 499 500 501 502 503 504 505 506 507 508 509 510 511 512 513 514 515 516 517 518 519 520 521 522 523 524 525 526 527 528 529 530 531 532 533 534 535 536 537 538 539 540 541 542 543 544 545 546 547 548 549 550 551 552 553 554 555 556 557 558 559 560 561 562 563 564 565 566 567 568 569 570 571 572 573 574 575 576 577 578 579 580 581 582 583 584 585 586 587 588 589 590 591 592 593 594 595 596 597 598 599 600 601 602 603 604 605 606 607 608 609 610 611 612 613 614 615 616 617 618 619 620 621 622 623 624 625 626 627 628 629 630 631 632 633 634 635 636 637 638 639 640 641 642 643 644 645 646 647 648 649 650 651 652 653 654 655 656 657 658 659 660 661 662 663 664 665 666 667 668 669 670 671 672 673 674 675 676 677 678 679 680 681 682 683 684 685 686 687 688 689 690 691 692 693 694 695 696 697 698 699 700 701 702 703 704 705 706 707 708 709 710 711 712 713 714 715 716 717 718 719 720 721 722 723 724 725 726 727 728 729 730 731 732 733 734 735 736 737 738 739 740 741 742 743 744 745 746 747 748 749 750 751 752 753 754 755 756 757 758 759 760 761 762 763 764 765 766 767 768 769 770 771 772 773 774 775 776 777 778 779 780 781 782 783 784 785 786 787 788 789 790 791 792 793 794 795 796 797 798 799 800 801 802 803 804 805 806 807 808 809 810 811 812 813 814 815 816 817 818 819 820 821 822 823 824 825 826 827 828 829 830 831 832 833 834 835 836 837 838 839 840 841 842 843 844 845 846 847 848 849 850 851 852 853 854 855 856 857 858 859 860 861 862 863 864 865 866 867 868 869 870 871 872 873 874 875 876 877 878 879 880 881 882 883 884 885 886 887 888 889 890 891 892 893 894 895 896 897 898 899 900 901 902 903 904 905 906 907 908 909 910 911 912 913 914 915 916 917 918 919 920 921 922 923 924 925 926 927 928 929 930 931 932 933 934 935 936 937 938 939 940 941 942 943 944 945 946 947 948 949 950 951 952 953 954 955 956 957 958 959 960 961 962 963 964 965 966 967 968 969 970 971 972 973 974 975 976 977 978 979 980 981 982 983 984 985 986 987 988 989 990 991 992 993 994 995 996 997 998 999 1000 1001 1002 1003 1004 1005 1006 1007 1008 1009 1010 1011 1012 1013 1014 1015 1016 1017 1018 1019 1020 1021 1022 1023 1024 1025 1026 1027 1028 1029 1030 1031 1032 1033 1034 1035 1036 1037 1038 1039 1040 1041 1042 1043 1044



TABLE B1.

**SUMMARY OF THE INCIDENCE OF NEOPLASMS IN MALE MICE  
ADMINISTERED N-NITROSODIPHENYLAMINE IN THE DIET**

	MATCHED CONTROL	LOW DOSE	HIGH DOSE
ANIMALS INITIALLY IN STUDY	20	50	50
ANIMALS MISSING			2
ANIMALS NECROPSIED	20	49	48
ANIMALS EXAMINED HISTOPATHOLOGICALLY	20	49	48
-----			
INTEGUMENTARY SYSTEM			
*SKIN	(20)	(49)	(48)
SEBACEOUS ADENOMA	1 (5%)		
HEMANGIOMA	1 (5%)		
-----			
RESPIRATORY SYSTEM			
#LUNG	(20)	(48)	(48)
CARCINOMA, NOS			1 (2%)
HEPATOCELLULAR CARCINOMA, METAST		1 (2%)	
ALVEOLAR/BRONCHIOLAR ADENOMA	4 (20%)	9 (19%)	7 (15%)
-----			
HEMATOPOIETIC SYSTEM			
*MULTIPLE ORGANS	(20)	(49)	(48)
MALIG.LYMPHOMA, HISTIOCYTIC TYPE			1 (2%)
GRANULOCYTIC LEUKEMIA	1 (5%)	1 (2%)	1 (2%)
*ABDOMINAL CAVITY	(20)	(49)	(48)
MALIG.LYMPHOMA, HISTIOCYTIC TYPE			1 (2%)
*BLOOD	(20)	(49)	(48)
LYMPHOCYTIC LEUKEMIA			1 (2%)
*SPLEEN	(20)	(49)	(48)
HEMANGIOMA			2 (4%)
HEMANGIOSARCOMA		2 (4%)	2 (4%)
*SMALL INTESTINE	(20)	(47)	(46)
MALIG.LYMPHOMA, UNDIFFER-TYPE		1 (2%)	
-----			
CIRCULATORY SYSTEM			
-----NONE-----			
-----			
# NUMBER OF ANIMALS WITH TISSUE EXAMINED MICROSCOPICALLY			
* NUMBER OF ANIMALS NECROPSIED			

TABLE B1. MALE MICE: NEOPLASMS (CONTINUED)

	MATCHED CONTROL	LOW DOSE	HIGH DOSE
DIGESTIVE SYSTEM			
#SALIVARY GLAND NEOPLASM, NOS, MALIGNANT	(20)	(48) 1 (2%)	(45)
#LIVER NEOPLASM, NOS, METASTATIC	(20)	(49) 1 (2%)	(48)
HEPATOCELLULAR ADENOMA	6 (30%)	11 (22%)	7 (15%)
HEPATOCELLULAR CARCINOMA		1 (2%)	
HEMANGIOMA		3 (6%)	
HEMANGIOSARCOMA		1 (2%)	
HEMANGIOSARCOMA, METASTATIC		2 (4%)	
#STOMACH PAPILLARY ADENOMA	(20) 1 (5%)	(48)	(46)
URINARY SYSTEM			
#KIDNEY TUBULAR-CELL ADENOMA	(20)	(49)	(48) 1 (2%)
#URINARY BLADDER PAPILLOMA, NOS	(18)	(49) 1 (2%)	(46)
TRANSITIONAL-CELL PAPILLOMA			1 (2%)
TRANSITIONAL-CELL CARCINOMA		1 (2%)	
ENDOCRINE SYSTEM			
#ADRENAL PHEOCHROMOCYTOMA	(18)	(49) 1 (2%)	(48)
REPRODUCTIVE SYSTEM			
NONE			
NERVOUS SYSTEM			
NONE			
SPECIAL SENSE ORGANS			
*EYE/LACRIMAL GLAND ADENOMA, NOS	(20)	(49) 1 (2%)	(48) 1 (2%)
# NUMBER OF ANIMALS WITH TISSUE EXAMINED MICROSCOPICALLY * NUMBER OF ANIMALS NECROPSIED			

TABLE B1. MALE MICE: NEOPLASMS (CONTINUED)

	MATCHED CONTROL	LOW DOSE	HIGH DOSE
CYSTADENOMA, NOS		1 (2%)	
MUSCULOSKELETAL SYSTEM			
NONE			
BODY CAVITIES			
NONE			
ALL OTHER SYSTEMS			
*MULTIPLE ORGANS	(20)	(49)	(48)
NEOPLASM, NOS, MALIGNANT		1 (2%)	
HEMANGIOMA			1 (2%)
HEMANGIOSARCOMA		1 (2%)	
ANIMAL DISPOSITION SUMMARY			
ANIMALS INITIALLY IN STUDY	20	50	50
NATURAL DEATH@	2	4	6
MORIBUND SACRIFICE			
SCHEDULED SACRIFICE			
ACCIDENTALLY KILLED			1
TERMINAL SACRIFICE	18	46	41
ANIMAL MISSING			2
@ INCLUDES AUTOLYZED ANIMALS			
* NUMBER OF ANIMALS WITH TISSUE EXAMINED MICROSCOPICALLY			
* NUMBER OF ANIMALS NECROPSIED			

**TABLE B1. MALE MICE: NEOPLASMS (CONTINUED)**

	MATCHED CONTROL	LOW DOSE	HIGH DOSE
TUMOR SUMMARY			
TOTAL ANIMALS WITH PRIMARY TUMORS*	10	29	22
TOTAL PRIMARY TUMORS	14	37	27
TOTAL ANIMALS WITH BENIGN TUMORS	9	23	18
TOTAL BENIGN TUMORS	13	27	20
TOTAL ANIMALS WITH MALIGNANT TUMORS	1	9	6
TOTAL MALIGNANT TUMORS	1	10	7
TOTAL ANIMALS WITH SECONDARY TUMORS#		4	
TOTAL SECONDARY TUMORS		4	
TOTAL ANIMALS WITH TUMORS UNCERTAIN- BENIGN OR MALIGNANT			
TOTAL UNCERTAIN TUMORS			
TOTAL ANIMALS WITH TUMORS UNCERTAIN- PRIMARY OR METASTATIC			
TOTAL UNCERTAIN TUMORS			
* PRIMARY TUMORS: ALL TUMORS EXCEPT SECONDARY TUMORS			
# SECONDARY TUMORS: METASTATIC TUMORS OR TUMORS INVASIVE INTO AN ADJACENT ORGAN			

TABLE B2.

**SUMMARY OF THE INCIDENCE OF NEOPLASMS IN FEMALE MICE  
ADMINISTERED N-NITROSODIPHENYLAMINE IN THE DIET**

	MATCHED CONTROL	LOW DOSE	HIGH DOSE
ANIMALS INITIALLY IN STUDY	20	50	50
ANIMALS MISSING			7
ANIMALS NECROPSIED	20	50	41
ANIMALS EXAMINED HISTOPATHOLOGICALLY	20	50	40
INTEGUMENTARY SYSTEM			
*SUBCUT TISSUE	(20)	(50)	(41)
BASAL-CELL TUMOR			1 (2%)
RHABDOMYOSARCOMA		1 (2%)	
RESPIRATORY SYSTEM			
#LUNG	(20)	(47)	(38)
CARCINOMA, NOS		1 (2%)	
ADENOCARCINOMA, NOS, METASTATIC		1 (2%)	
ALVEOLAR/BRONCHIOLAR ADENOMA	3 (15%)	9 (19%)	5 (13%)
ALVEOLAR/BRONCHIOLAR CARCINOMA		2 (4%)	
OSTEOSARCOMA, METASTATIC	1 (5%)		
HEMATOPOIETIC SYSTEM			
*MULTIPLE ORGANS	(20)	(50)	(41)
MALIG.LYMPHOMA, UNDIFFER-TYPE		1 (2%)	
MALIG.LYMPHOMA, HISTIOCYTIC TYPE		1 (2%)	
PLASMA-CELL TUMOR		1 (2%)	
LYMPHOCYTIC LEUKEMIA		1 (2%)	
GRANULOCYTIC LEUKEMIA			1 (2%)
*ABDOMINAL CAVITY	(20)	(50)	(41)
MALIG.LYMPHOMA, HISTIOCYTIC TYPE			1 (2%)
#SPLEEN	(20)	(48)	(38)
SARCOMA, NOS	1 (5%)		
HEMANGIOMA			1 (3%)
MALIG.LYMPHOMA, UNDIFFER-TYPE			1 (3%)
#SMALL INTESTINE	(19)	(48)	(39)
MALIG.LYMPHOMA, UNDIFFER-TYPE		1 (2%)	
* NUMBER OF ANIMALS WITH TISSUE EXAMINED MICROSCOPICALLY			
* NUMBER OF ANIMALS NECROPSIED			



TABLE B2. FEMALE MICE: NEOPLASMS (CONTINUED)

	MATCHED CONTROL	LOW DOSE	HIGH DOSE
*MESENTERY MALIG. LYMPHOMA, UNDIFFER-TYPE	(20)	(50)	(41) 1 (2%)
#KIDNEY MALIG. LYMPHOMA, HISTIOCYTIC TYPE	(20) 1 (5%)	(49)	(38)
CIRCULATORY SYSTEM			
NONE			
DIGESTIVE SYSTEM			
#LIVER	(20)	(49)	(38)
HEPATOCELLULAR ADENOMA	3 (15%)	7 (14%)	4 (11%)
HEPATOCELLULAR CARCINOMA	1 (5%)		
HEMANGIOMA			1 (3%)
URINARY SYSTEM			
#URINARY BLADDER	(18)	(47)	(38)
TRANSITIONAL-CELL CARCINOMA		1 (2%)	
#U. BLADDER/SUBMUCOSA	(18)	(47)	(38)
HEMANGIOMA		1 (2%)	
ENDOCRINE SYSTEM			
#PITUITARY	(14)	(37)	(26)
CHROMOPHOBE ADENOMA		1 (3%)	
#ADRENAL	(20)	(45)	(37)
CORTICAL ADENOMA	1 (5%)		
REPRODUCTIVE SYSTEM			
#UTERUS	(20)	(50)	(37)
LEIOMYOMA		1 (2%)	
HEMANGIOMA			1 (3%)
#OVARY/FOLLICLE	(20)	(48)	(38)
ADENOMA, NOS			1 (3%)

# NUMBER OF ANIMALS WITH TISSUE EXAMINED MICROSCOPICALLY

\* NUMBER OF ANIMALS NECROPSIED

**TABLE B2. FEMALE MICE: NEOPLASMS (CONTINUED)**

	MATCHED CONTROL	LOW DOSE	HIGH DOSE
NERVOUS SYSTEM			
NONE			
SPECIAL SENSE ORGANS			
*EYE/LACRIMAL GLAND ADENOMA, NOS	(20)	(50)	(41) 1 (2%)
MUSCULOSKELETAL SYSTEM			
*SKULL OSTEOSARCOMA	(20) 1 (5%)	(50)	(41)
*SKELETAL MUSCLE RHABDOMYOSARCOMA	(20)	(50) 1 (2%)	(41)
BODY CAVITIES			
NONE			
ALL OTHER SYSTEMS			
NONE			
ANIMAL DISPOSITION SUMMARY			
ANIMALS INITIALLY IN STUDY	20	50	50
NATURAL DEATH@	4	7	11
MORIBUND SACRIFICE		1	1
SCHEDULED SACRIFICE			
ACCIDENTALLY KILLED			
TERMINAL SACRIFICE	16	42	31
ANIMAL MISSING			7
@ INCLUDES AUTOLYZED ANIMALS			
* NUMBER OF ANIMALS WITH TISSUE EXAMINED MICROSCOPICALLY			
* NUMBER OF ANIMALS NECROPSIED			

**TABLE B2. FEMALE MICE: NEOPLASMS (CONTINUED)**

	MATCHED CONTROL	LOW DOSE	HIGH DOSE
<b>TUMOR SUMMARY</b>			
TOTAL ANIMALS WITH PRIMARY TUMORS*	9	23	15
TOTAL PRIMARY TUMORS	11	30	19
TOTAL ANIMALS WITH BENIGN TUMORS	7	17	13
TOTAL BENIGN TUMORS	7	19	15
TOTAL ANIMALS WITH MALIGNANT TUMORS	3	10	4
TOTAL MALIGNANT TUMORS	4	10	4
TOTAL ANIMALS WITH SECONDARY TUMORS#	1	1	
TOTAL SECONDARY TUMORS	1	1	
TOTAL ANIMALS WITH TUMORS UNCERTAIN- BENIGN OR MALIGNANT		1	
TOTAL UNCERTAIN TUMORS		1	
TOTAL ANIMALS WITH TUMORS UNCERTAIN- PRIMARY OR METASTATIC			
TOTAL UNCERTAIN TUMORS			

\* PRIMARY TUMORS: ALL TUMORS EXCEPT SECONDARY TUMORS

# SECONDARY TUMORS: METASTATIC TUMORS OR TUMORS INVASIVE INTO AN ADJACENT ORGAN

## APPENDIX C

### SUMMARY OF THE INCIDENCE OF NONNEOPLASTIC LESIONS IN RATS ADMINISTERED N-NITROSODIPHENYLAMINE IN THE DIET

[illegible]

TABLE C1.

**SUMMARY OF THE INCIDENCE OF NONNEOPLASTIC LESIONS IN MALE RATS  
ADMINISTERED N-NITROSODIPHENYLAMINE IN THE DIET**

	MATCHED CONTROL	LOW DOSE	HIGH DOSE
ANIMALS INITIALLY IN STUDY	20	50	50
ANIMALS NECROPSIED	20	50	50
ANIMALS EXAMINED HISTOPATHOLOGICALLY	20	50	50
-----			
INTEGUMENTARY SYSTEM			
*SKIN	(20)	(50)	(50)
HYPERKERATOSIS		1 (2%)	
*SUBCUT TISSUE	(20)	(50)	(50)
INFLAMMATION, ACUTE			1 (2%)
INFLAMMATION, CHRONIC			1 (2%)
-----			
RESPIRATORY SYSTEM			
#LUNG	(20)	(49)	(49)
HYPERPLASIA, ADENOMATOUS	1 (5%)		1 (2%)
-----			
HEMATOPOIETIC SYSTEM			
#MANDIBULAR L. NODE	(19)	(50)	(50)
HYPERPLASIA, NOS		1 (2%)	
-----			
CIRCULATORY SYSTEM			
#MYOCARDIUM	(20)	(50)	(49)
FIBROSIS			1 (2%)
-----			
DIGESTIVE SYSTEM			
#LIVER	(20)	(50)	(50)
DEGENERATION, NOS	2 (10%)	5 (10%)	1 (2%)
HYPERPLASIA, NODULAR			1 (2%)
HYPERPLASIA, FOCAL	1 (5%)		1 (2%)
#LIVER/CENTRIOLOBULAR	(20)	(50)	(50)
DEGENERATION, NOS	1 (5%)		
-----			
# NUMBER OF ANIMALS WITH TISSUE EXAMINED MICROSCOPICALLY			
* NUMBER OF ANIMALS NECROPSIED			



**TABLE C1. MALE RATS: NONNEOPLASTIC LESIONS (CONTINUED)**

	MATCHED CONTROL	LOW DOSE	HIGH DOSE
NECROSIS, NOS	1 (5%)		
#BILE DUCT HYPERPLASIA, NOS	(20)	(50) 1 (2%)	(50)
#PANCREATIC ACINUS HYPERPLASIA, NODULAR	(18) 1 (6%)	(45)	(36)
URINARY SYSTEM			
#URINARY BLADDER HYPERPLASIA, EPITHELIAL METAPLASIA, SQUAMOUS	(19)	(46) 2 (4%)	(45) 6 (13%) 1 (2%)
ENDOCRINE SYSTEM			
#PITUITARY ANGIECTASIS	(18)	(47)	(47) 1 (2%)
#ADRENAL HEMORRHAGIC CYST	(19) 1 (5%)	(46)	(49)
#ADRENAL CORTEX HYPERPLASIA, FOCAL	(19)	(46) 1 (2%)	(49)
#ADRENAL MEDULLA HYPERPLASIA, NOS HYPERPLASIA, FOCAL	(19) 1 (5%)	(46) 1 (2%)	(49) 2 (4%)
#THYROID CYSTIC FOLLICLES PIGMENTATION, NOS HYPERPLASIA, C-CELL	(19) 1 (5%)	(49) 2 (4%) 1 (2%) 1 (2%)	(48) 1 (2%)
#PANCREATIC ISLETS HYPERPLASIA, NOS	(18)	(45) 1 (2%)	(36)
REPRODUCTIVE SYSTEM			
*MAMMARY GLAND CYST, NOS METAPLASIA, SQUAMOUS	(20)	(50) 1 (2%) 1 (2%)	(50)

# NUMBER OF ANIMALS WITH TISSUE EXAMINED MICROSCOPICALLY

\* NUMBER OF ANIMALS NECROPSIED

**TABLE C1. MALE RATS: NONNEOPLASTIC LESIONS (CONTINUED)**

	MATCHED CONTROL	LOW DOSE	HIGH DOSE
#TESTIS	(19)	(50)	(49)
HYPERPLASIA, INTERSTITIAL CELL	1 (5%)	1 (2%)	
NERVOUS SYSTEM			
NONE			
SPECIAL SENSE ORGANS			
NONE			
MUSCULOSKELETAL SYSTEM			
NONE			
BODY CAVITIES			
NONE			
ALL OTHER SYSTEMS			
NONE			
SPECIAL MORPHOLOGY SUMMARY			
NO LESION REPORTED	1	1	3
# NUMBER OF ANIMALS WITH TISSUE EXAMINED MICROSCOPICALLY			
* NUMBER OF ANIMALS NECROPSIED			

TABLE C2.

**SUMMARY OF THE INCIDENCE OF NONNEOPLASTIC LESIONS IN FEMALE RATS  
ADMINISTERED N-NITROSODIPHENYLAMINE IN THE DIET**

	MATCHED CONTROL	LOW DOSE	HIGH DOSE
ANIMALS INITIALLY IN STUDY	20	50	50
ANIMALS MISSING			1
ANIMALS NECROPSIED	20	50	49
ANIMALS EXAMINED HISTOPATHOLOGICALLY	20	50	49
-----			
INTEGUMENTARY SYSTEM			
*SUBCUT TISSUE	(20)	(50)	(49)
NECROSIS, NOS			1 (2%)
-----			
RESPIRATORY SYSTEM			
#LUNG	(19)	(50)	(46)
PNEUMONIA, CHRONIC MURINE	1 (5%)		
HYPERPLASIA, ADENOMATOUS	1 (5%)		
-----			
HEMATOPOIETIC SYSTEM			
*BLOOD	(20)	(50)	(49)
POLYCHROMASIA		2 (4%)	
#SPLEEN	(20)	(48)	(49)
HEMATOPOIESIS		1 (2%)	
#THYMUS	(19)	(48)	(23)
HYPERTROPHY, NOS		1 (2%)	
-----			
CIRCULATORY SYSTEM			
NONE			
-----			
DIGESTIVE SYSTEM			
*LIVER	(20)	(50)	(49)
INFLAMMATION, ACUTE			1 (2%)
INFLAMMATION, FOCAL GRANULOMATOUS		1 (2%)	
-----			
# NUMBER OF ANIMALS WITH TISSUE EXAMINED MICROSCOPICALLY			
* NUMBER OF ANIMALS NECROPSIED			

**TABLE C2. FEMALE RATS: NONNEOPLASTIC LESIONS (CONTINUED)**

	MATCHED CONTROL	LOW DOSE	HIGH DOSE
DEGENERATION, NOS	1 (5%)	2 (4%)	
NECROSIS, COAGULATIVE			1 (2%)
HYPERPLASIA, NODULAR	1 (5%)		
HYPERPLASTIC NODULE			1 (2%)
HYPERPLASIA, FOCAL		3 (6%)	
URINARY SYSTEM			
#URINARY BLADDER	(18)	(48)	(49)
FIBROSIS			1 (2%)
HYPERPLASIA, EPITHELIAL		4 (8%)	7 (14%)
METAPLASIA, SQUAMOUS			2 (4%)
#U. BLADDER/SUBMUCOSA	(18)	(48)	(49)
INFLAMMATION, CHRONIC		1 (2%)	
ENDOCRINE SYSTEM			
#PITUITARY	(19)	(49)	(48)
HEMORRHAGE		1 (2%)	
HEMORRHAGIC CYST		1 (2%)	
ANGIECTASIS		1 (2%)	
#ADRENAL	(19)	(50)	(48)
LIPOIDOSIS		1 (2%)	
ANGIECTASIS		1 (2%)	
#ADRENAL CORTEX	(19)	(50)	(48)
HYPERPLASIA, FOCAL		1 (2%)	
#THYROID	(20)	(49)	(48)
CYSTIC FOLLICLES		3 (6%)	
HYPERPLASIA, C-CELL			1 (2%)
REPRODUCTIVE SYSTEM			
*MAMMARY GLAND	(20)	(50)	(49)
DILATATION/DUCTS	3 (15%)	3 (6%)	5 (10%)
FIBROSIS			1 (2%)
LACTATION			2 (4%)
#UTERUS	(20)	(50)	(46)
HYDROMETRA		1 (2%)	
# NUMBER OF ANIMALS WITH TISSUE EXAMINED MICROSCOPICALLY			
* NUMBER OF ANIMALS NECROPSIED			

**TABLE C2. FEMALE RATS: NONNEOPLASTIC LESIONS (CONTINUED)**

	MATCHED CONTROL	LOW DOSE	HIGH DOSE
THROMBUS, ORGANIZED	1 (5%)		
FIBROSIS		1 (2%)	
NECROSIS, NOS	1 (5%)		
POLYP	3 (15%)	5 (10%)	1 (2%)
*UTERUS/ENDOMETRIUM	(20)	(50)	(46)
INFLAMMATION, NECROTIZING			1 (2%)
HYPERPLASIA, CYSTIC		1 (2%)	1 (2%)
*OVARY	(20)	(50)	(49)
FOLLICULAR CYST, NOS		1 (2%)	1 (2%)
NERVOUS SYSTEM			
NONE			
SPECIAL SENSE ORGANS			
NONE			
MUSCULOSKELETAL SYSTEM			
NONE			
BODY CAVITIES			
NONE			
ALL OTHER SYSTEMS			
*MULTIPLE ORGANS	(20)	(50)	(49)
POSTMORTEM CHANGE		1 (2%)	1 (2%)
SPECIAL MORPHOLOGY SUMMARY			
NO LESION REPORTED	6	13	
ANIMAL MISSING/NO NECROPSY			1
* NUMBER OF ANIMALS WITH TISSUE EXAMINED MICROSCOPICALLY			
* NUMBER OF ANIMALS NECROPSIED			

APPENDIX D

SUMMARY OF THE INCIDENCE OF NONNEOPLASTIC LESIONS IN MICE  
ADMINISTERED N-NITROSODIPHENYLAMINE IN THE DIET



100  
101  
102  
103  
104  
105  
106  
107  
108  
109  
110  
111  
112  
113  
114  
115  
116  
117  
118  
119  
120  
121  
122  
123  
124  
125  
126  
127  
128  
129  
130  
131  
132  
133  
134  
135  
136  
137  
138  
139  
140  
141  
142  
143  
144  
145  
146  
147  
148  
149  
150  
151  
152  
153  
154  
155  
156  
157  
158  
159  
160  
161  
162  
163  
164  
165  
166  
167  
168  
169  
170  
171  
172  
173  
174  
175  
176  
177  
178  
179  
180  
181  
182  
183  
184  
185  
186  
187  
188  
189  
190  
191  
192  
193  
194  
195  
196  
197  
198  
199  
200

TABLE D1.

**SUMMARY OF THE INCIDENCE OF NONNEOPLASTIC LESIONS IN MALE MICE  
ADMINISTERED N-NITROSODIPHENYLAMINE IN THE DIET**

	MATCHED CONTROL	LOW DOSE	HIGH DOSE
ANIMALS INITIALLY IN STUDY	20	50	50
ANIMALS MISSING			2
ANIMALS NECROPSIED	20	49	48
ANIMALS EXAMINED HISTOPATHOLOGICALLY	20	49	48
INTEGUMENTARY SYSTEM			
*SUBCUT TISSUE	(20)	(49)	(48)
CYST, NOS			1 (2%)
NECROSIS, FAT			1 (2%)
RESPIRATORY SYSTEM			
NONE			
HEMATOPOIETIC SYSTEM			
*BONE MARROW	(18)	(49)	(48)
HYPERPLASIA, HEMATOPOIETIC		1 (2%)	
*SPLEEN	(20)	(49)	(48)
HEMATOPOIESIS			1 (2%)
*LYMPH NODE	(20)	(48)	(46)
HYPERPLASIA, NOS	1 (5%)		
*MESENTERIC L. NODE	(20)	(48)	(46)
HEMORRHAGE		1 (2%)	
HYPERPLASIA, NOS			1 (2%)
CIRCULATORY SYSTEM			
NONE			
DIGESTIVE SYSTEM			
*LIVER	(20)	(49)	(48)
THROMBOSIS, NOS		2 (4%)	
* NUMBER OF ANIMALS WITH TISSUE EXAMINED MICROSCOPICALLY			
* NUMBER OF ANIMALS NECROPSIED			

TABLE D1. MALE MICE: NONNEOPLASTIC LESIONS (CONTINUED)

	MATCHED CONTROL	LOW DOSE	HIGH DOSE
HEMORRHAGE		1 (2%)	
DEGENERATION, NOS			1 (2%)
NECROSIS, NOS		2 (4%)	
METAMORPHOSIS FATTY	2 (10%)	3 (6%)	1 (2%)
HYPERPLASIA, NODULAR	1 (5%)	1 (2%)	
HYPERPLASIA, FOCAL	1 (5%)		3 (6%)
#HEPATIC CAPSULE FIBROSIS	(20)	(49) 1 (2%)	(48)
#SMALL INTESTINE HYPERPLASIA, ADENOMATOUS	(20)	(47) 1 (2%)	(46)
#DUODENUM HYPERPLASIA, NOS	(20)	(47)	(46) 1 (2%)
URINARY SYSTEM			
#KIDNEY	(20)	(49)	(48)
PYELONEPHRITIS, ACUTE			1 (2%)
INFLAMMATION, CHRONIC		9 (18%)	5 (10%)
PERIVASCULAR CUFFING		1 (2%)	
#URINARY BLADDER	(18)	(49)	(46)
INFLAMMATION, ACUTE			1 (2%)
INFLAMMATION, CHRONIC		2 (4%)	
CYTOLOGIC DEGENERATION			1 (2%)
HYPERPLASIA, EPITHELIAL		2 (4%)	7 (15%)
#U. BLADDER/SUBMUCOSA	(18)	(49)	(46)
INFLAMMATION, NOS		1 (2%)	
INFLAMMATION, CHRONIC		12 (24%)	31 (67%)
HYPERPLASIA, LYMPHOID			1 (2%)
ENDOCRINE SYSTEM			
#THYROID	(19)	(47)	(45)
CYSTIC FOLLICLES			1 (2%)
HEMORRHAGE			1 (2%)
HYPERPLASIA, FOLLICULAR-CELL		1 (2%)	
#PANCREATIC ISLETS	(17)	(45)	(43)
HYPERPLASIA, NOS	1 (6%)		

# NUMBER OF ANIMALS WITH TISSUE EXAMINED MICROSCOPICALLY

\* NUMBER OF ANIMALS NECROPSIED

TABLE D1. MALE MICE: NONNEOPLASTIC LESIONS (CONTINUED)

	MATCHED CONTROL	LOW DOSE	HIGH DOSE
REPRODUCTIVE SYSTEM			
*SEMINAL VESICLE DISTENTION	(20) 1 (5%)	(49)	(48)
NERVOUS SYSTEM			
*MIDBRAIN HEMORRHAGE	(19)	(49)	(48) 1 (2%)
*MEDULLA OBLONGATA HEMORRHAGE	(19)	(49)	(48) 1 (2%)
SPECIAL SENSE ORGANS			
NONE			
MUSCULOSKELETAL SYSTEM			
NONE			
BODY CAVITIES			
*ABDOMINAL CAVITY NECROSIS, FAT	(20)	(49) 1 (2%)	(48) 1 (2%)
*MESENTERY NECROSIS, FAT	(20) 1 (5%)	(49)	(48)
ALL OTHER SYSTEMS			
NONE			
SPECIAL MORPHOLOGY SUMMARY			
NO LESION REPORTED	6	11	4
ANIMAL MISSING/NO NECROPSY			2
* NUMBER OF ANIMALS WITH TISSUE EXAMINED MICROSCOPICALLY			
* NUMBER OF ANIMALS NECROPSIED			

**TABLE D1. MALE MICE: NONNEOPLASTIC LESIONS (CONTINUED)**

	MATCHED CONTROL	LOW DOSE	HIGH DOSE
AUTO/NECROPSY/HISTO PERF			1
AUTOLYSIS/NO NECROPSY		1	
* NUMBER OF ANIMALS WITH TISSUE EXAMINED MICROSCOPICALLY			
* NUMBER OF ANIMALS NECROPSIED			

TABLE D2.

**SUMMARY OF THE INCIDENCE OF NONNEOPLASTIC LESIONS IN FEMALE MICE  
ADMINISTERED N-NITROSODIPHENYLAMINE IN THE DIET**

	MATCHED CONTROL	LOW DOSE	HIGH DOSE
ANIMALS INITIALLY IN STUDY	20	50	50
ANIMALS MISSING			7
ANIMALS NECROPSIED	20	50	41
ANIMALS EXAMINED HISTOPATHOLOGICALLY	20	50	40
INTEGUMENTARY SYSTEM			
*SUBCUT TISSUE	(20)	(50)	(41)
THROMBOSIS, NOS	1 (5%)		
HEMORRHAGE	2 (10%)		
INFLAMMATION, ACUTE/CHRONIC	2 (10%)		
RESPIRATORY SYSTEM			
*LUNG	(20)	(47)	(38)
NECROSIS, CENTRAL		1 (2%)	
HYPERPLASIA, ADENOMATOUS			1 (3%)
HEMATOPOIETIC SYSTEM			
*SPLEEN	(20)	(48)	(38)
THROMBOSIS, NOS			1 (3%)
*MESENTERIC L. NODE	(20)	(47)	(38)
HEMORRHAGE	1 (5%)		
HYPERPLASIA, NOS	1 (5%)		
*THYMUS	(17)	(38)	(37)
CONGESTION, NOS	1 (6%)		
HEMORRHAGE	1 (6%)		
HYPERPLASIA, NOS	1 (6%)		
CIRCULATORY SYSTEM			
NONE			
DIGESTIVE SYSTEM			
*LIVER	(20)	(49)	(38)
THROMBOSIS, NOS			1 (3%)
* NUMBER OF ANIMALS WITH TISSUE EXAMINED MICROSCOPICALLY			
* NUMBER OF ANIMALS NECROPSIED			



**TABLE D2. FEMALE MICE: NONNEOPLASTIC LESIONS (CONTINUED)**

	MATCHED CONTROL	LOW DOSE	HIGH DOSE
INFLAMMATION, ACUTE FOCAL			1 (3%)
DEGENERATION, NOS	1 (5%)		1 (3%)
NECROSIS, NOS		1 (2%)	
NECROSIS, COAGULATIVE			1 (3%)
INFARCT, NOS		1 (2%)	
#LIVER/CENTRILOBULAR NECROSIS, NOS	(20)	(49)	(38) 1 (3%)
#PANCREAS	(18)	(42)	(35)
INFLAMMATION, ACUTE			1 (3%)
INFLAMMATION, ACUTE/CHRONIC	1 (6%)		
INFLAMMATION, CHRONIC FOCAL			1 (3%)
#STOMACH	(19)	(49)	(39)
INFLAMMATION, ACUTE			1 (3%)
INFLAMMATION, ACUTE FOCAL	1 (5%)		
INFLAMMATION, ACUTE/CHRONIC			1 (3%)
HYPERPLASIA, EPITHELIAL			1 (3%)
#GASTRIC SUBMUCOSA	(19)	(49)	(39)
INFLAMMATION, CHRONIC			1 (3%)
#SMALL INTESTINE	(19)	(48)	(39)
HYPERPLASIA, ADENOMATOUS		1 (2%)	
<b>URINARY SYSTEM</b>			
#KIDNEY	(20)	(49)	(38)
INFLAMMATION, CHRONIC	5 (25%)		14 (37%)
PERIVASCULAR CUFFING		5 (10%)	
#URINARY BLADDER	(18)	(47)	(38)
EDEMA, NOS			1 (3%)
LYMPHOCYTIC INFLAMMATORY INFILTR		1 (2%)	
INFLAMMATION, CHRONIC			5 (13%)
PERIVASCULITIS			2 (5%)
DEGENERATION, NOS		1 (2%)	
NECROSIS, FIBRINOID			1 (3%)
HYPERPLASIA, EPITHELIAL		3 (6%)	5 (13%)
#U. BLADDER/SUBMUCOSA	(18)	(47)	(38)
EDEMA, NOS		1 (2%)	
LYMPHOCYTIC INFLAMMATORY INFILTR	1 (6%)		

# NUMBER OF ANIMALS WITH TISSUE EXAMINED MICROSCOPICALLY

\* NUMBER OF ANIMALS NECROPSIED

TABLE D2. FEMALE MICE: NONNEOPLASTIC LESIONS (CONTINUED)

	MATCHED CONTROL	LOW DOSE	HIGH DOSE
INFLAMMATION, CHRONIC		31 (66%)	30 (79%)
HYPERPLASIA, EPITHELIAL			1 (3%)
ENDOCRINE SYSTEM			
#THYROID	(19)	(47)	(37)
CYSTIC FOLLICLES			1 (3%)
REPRODUCTIVE SYSTEM			
#UTERUS/ENDOMETRIUM	(20)	(50)	(37)
CYST, NOS		5 (10%)	
INFLAMMATION, NOS	1 (5%)		
INFLAMMATION, ACUTE		1 (2%)	
HYPERPLASIA, NOS		1 (2%)	
HYPERPLASIA, CYSTIC	6 (30%)	9 (18%)	2 (5%)
#OVARY	(20)	(48)	(38)
FOLLICULAR CYST, NOS		2 (4%)	3 (8%)
PAROVARIAN CYST	1 (5%)		
INFLAMMATION, ACUTE/CHRONIC	1 (5%)		
HYPERPLASIA, GRANULOSA-CELL		1 (2%)	
NERVOUS SYSTEM			
NONE			
SPECIAL SENSE ORGANS			
NONE			
MUSCULOSKELETAL SYSTEM			
NONE			
BODY CAVITIES			
NONE			
ALL OTHER SYSTEMS			
NONE			
# NUMBER OF ANIMALS WITH TISSUE EXAMINED MICROSCOPICALLY			
* NUMBER OF ANIMALS NECROPSIED			

**TABLE D2. FEMALE MICE: NONNEOPLASTIC LESIONS (CONTINUED)**

	MATCHED CONTROL	LOW DOSE	HIGH DOSE
SPECIAL MORPHOLOGY SUMMARY			
NO LESION REPORTED	4	6	1
ANIMAL MISSING/NO NECROPSY			7
NECROPSY PERF/NO HISTO PERFORMED			1
AUTO/NECROPSY/HISTO PERF		2	1
AUTOLYSIS/NO NECROPSY			2
# NUMBER OF ANIMALS WITH TISSUE EXAMINED MICROSCOPICALLY			
* NUMBER OF ANIMALS NECROPSIED			

APPENDIX E

ANALYSES OF THE INCIDENCE OF PRIMARY TUMORS IN RATS  
ADMINISTERED N-NITROSODIPHENYLAMINE IN THE DIET

11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65  
66  
67  
68  
69  
70  
71  
72  
73  
74  
75  
76  
77  
78  
79  
80  
81  
82  
83  
84  
85  
86  
87  
88  
89  
90  
91  
92  
93  
94  
95  
96  
97  
98  
99  
100  
101  
102  
103  
104  
105  
106  
107  
108  
109  
110  
111  
112  
113  
114  
115  
116  
117  
118  
119  
120  
121  
122  
123  
124  
125  
126  
127  
128  
129  
130  
131  
132  
133  
134  
135  
136  
137  
138  
139  
140  
141  
142  
143  
144  
145  
146  
147  
148  
149  
150  
151  
152  
153  
154  
155  
156  
157  
158  
159  
160  
161  
162  
163  
164  
165  
166  
167  
168  
169  
170  
171  
172  
173  
174  
175  
176  
177  
178  
179  
180  
181  
182  
183  
184  
185  
186  
187  
188  
189  
190  
191  
192  
193  
194  
195  
196  
197  
198  
199  
200  
201  
202  
203  
204  
205  
206  
207  
208  
209  
210  
211  
212  
213  
214  
215  
216  
217  
218  
219  
220  
221  
222  
223  
224  
225  
226  
227  
228  
229  
230  
231  
232  
233  
234  
235  
236  
237  
238  
239  
240  
241  
242  
243  
244  
245  
246  
247  
248  
249  
250  
251  
252  
253  
254  
255  
256  
257  
258  
259  
260  
261  
262  
263  
264  
265  
266  
267  
268  
269  
270  
271  
272  
273  
274  
275  
276  
277  
278  
279  
280  
281  
282  
283  
284  
285  
286  
287  
288  
289  
290  
291  
292  
293  
294  
295  
296  
297  
298  
299  
300  
301  
302  
303  
304  
305  
306  
307  
308  
309  
310  
311  
312  
313  
314  
315  
316  
317  
318  
319  
320  
321  
322  
323  
324  
325  
326  
327  
328  
329  
330  
331  
332  
333  
334  
335  
336  
337  
338  
339  
340  
341  
342  
343  
344  
345  
346  
347  
348  
349  
350  
351  
352  
353  
354  
355  
356  
357  
358  
359  
360  
361  
362  
363  
364  
365  
366  
367  
368  
369  
370  
371  
372  
373  
374  
375  
376  
377  
378  
379  
380  
381  
382  
383  
384  
385  
386  
387  
388  
389  
390  
391  
392  
393  
394  
395  
396  
397  
398  
399  
400  
401  
402  
403  
404  
405  
406  
407  
408  
409  
410  
411  
412  
413  
414  
415  
416  
417  
418  
419  
420  
421  
422  
423  
424  
425  
426  
427  
428  
429  
430  
431  
432  
433  
434  
435  
436  
437  
438  
439  
440  
441  
442  
443  
444  
445  
446  
447  
448  
449  
450  
451  
452  
453  
454  
455  
456  
457  
458  
459  
460  
461  
462  
463  
464  
465  
466  
467  
468  
469  
470  
471  
472  
473  
474  
475  
476  
477  
478  
479  
480  
481  
482  
483  
484  
485  
486  
487  
488  
489  
490  
491  
492  
493  
494  
495  
496  
497  
498  
499  
500  
501  
502  
503  
504  
505  
506  
507  
508  
509  
510  
511  
512  
513  
514  
515  
516  
517  
518  
519  
520  
521  
522  
523  
524  
525  
526  
527  
528  
529  
530  
531  
532  
533  
534  
535  
536  
537  
538  
539  
540  
541  
542  
543  
544  
545  
546  
547  
548  
549  
550  
551  
552  
553  
554  
555  
556  
557  
558  
559  
560  
561  
562  
563  
564  
565  
566  
567  
568  
569  
570  
571  
572  
573  
574  
575  
576  
577  
578  
579  
580  
581  
582  
583  
584  
585  
586  
587  
588  
589  
590  
591  
592  
593  
594  
595  
596  
597  
598  
599  
600  
601  
602  
603  
604  
605  
606  
607  
608  
609  
610  
611  
612  
613  
614  
615  
616  
617  
618  
619  
620  
621  
622  
623  
624  
625  
626  
627  
628  
629  
630  
631  
632  
633  
634  
635  
636  
637  
638  
639  
640  
641  
642  
643  
644  
645  
646  
647  
648  
649  
650  
651  
652  
653  
654  
655  
656  
657  
658  
659  
660  
661  
662  
663  
664  
665  
666  
667  
668  
669  
670  
671  
672  
673  
674  
675  
676  
677  
678  
679  
680  
681  
682  
683  
684  
685  
686  
687  
688  
689  
690  
691  
692  
693  
694  
695  
696  
697  
698  
699  
700  
701  
702  
703  
704  
705  
706  
707  
708  
709  
710  
711  
712  
713  
714  
715  
716  
717  
718  
719  
720  
721  
722  
723  
724  
725  
726  
727  
728  
729  
730  
731  
732  
733  
734  
735  
736  
737  
738  
739  
740  
741  
742  
743  
744  
745  
746  
747  
748  
749  
750  
751  
752  
753  
754  
755  
756  
757  
758  
759  
760  
761  
762  
763  
764  
765  
766  
767  
768  
769  
770  
771  
772  
773  
774  
775  
776  
777  
778  
779  
780  
781  
782  
783  
784  
785  
786  
787  
788  
789  
790  
791  
792  
793  
794  
795  
796  
797  
798  
799  
800  
801  
802  
803  
804  
805  
806  
807  
808  
809  
810  
811  
812  
813  
814  
815  
816  
817  
818  
819  
820  
821  
822  
823  
824  
825  
826  
827  
828  
829  
830  
831  
832  
833  
834  
835  
836  
837  
838  
839  
840  
841  
842  
843  
844  
845  
846  
847  
848  
849  
850  
851  
852  
853  
854  
855  
856  
857  
858  
859  
860  
861  
862  
863  
864  
865  
866  
867  
868  
869  
870  
871  
872  
873  
874  
875  
876  
877  
878  
879  
880  
881  
882  
883  
884  
885  
886  
887  
888  
889  
890  
891  
892  
893  
894  
895  
896  
897  
898  
899  
900  
901  
902  
903  
904  
905  
906  
907  
908  
909  
910  
911  
912  
913  
914  
915  
916  
917  
918  
919  
920  
921  
922  
923  
924  
925  
926  
927  
928  
929  
930  
931  
932  
933  
934  
935  
936  
937  
938  
939  
940  
941  
942  
943  
944  
945  
946  
947  
948  
949  
950  
951  
952  
953  
954  
955  
956  
957  
958  
959  
960  
961  
962  
963  
964  
965  
966  
967  
968  
969  
970  
971  
972  
973  
974  
975  
976  
977  
978  
979  
980  
981  
982  
983  
984  
985  
986  
987  
988  
989  
990  
991  
992  
993  
994  
995  
996  
997  
998  
999  
1000

Table E1. Analyses of the Incidence of Primary Tumors in Male Rats  
Administered N-Nitrosodiphenylamine in the Diet (a)

<u>Topography: Morphology</u>	<u>Matched Control</u>	<u>Low Dose</u>	<u>High Dose</u>
Integumentary System: Basal-cell Tumor (b)	1/20 (5)	5/50 (10)	3/50 (6)
P Values (c,d)	N.S.	N.S.	N.S.
Relative Risk (f)			
Lower Limit		2.000	1.200
Upper Limit		0.249	0.106
		92.596	61.724
Weeks to First Observed Tumor	100	81	61
Integumentary System: Fibroma (b)	1/20 (5)	1/50 (2)	10/50 (20)
P Values (c,d)	P = 0.003	N.S.	N.S.
Relative Risk (f)			
Lower Limit		0.400	4.000
Upper Limit		0.005	0.642
		30.802	169.457
Weeks to First Observed Tumor	100	100	100



Table El. Analyses of the Incidence of Primary Tumors in Male Rats  
Administered N-Nitrosodiphenylamine in the Diet (a)

(continued)				
<u>Topography: Morphology</u>	<u>Matched Control</u>	<u>Low Dose</u>	<u>High Dose</u>	
Hematopoietic System: Leukemia (b)	3/20 (15)	2/50 (4)	2/50 (4)	
P Values (c,d)	N.S.	N.S.	N.S.	
Relative Risk (f)				
Lower Limit		0.267	0.267	
Upper Limit		0.024	0.024	
		2.190	2.190	
Weeks to First Observed Tumor	88	95	93	
<hr/>				
Urinary Bladder: Transitional-cell Carcinoma (b)	0/19 (0)	0/46 (0)	16/45 (36)	
P Values (c,d)	P less than 0.001	--	P = 0.001	
Relative Risk (f)		--	Infinite	
Lower Limit		--	2.239	
Upper Limit		--	Infinite	
Weeks to First Observed Tumor	--	--	97	

Table El. Analyses of the Incidence of Primary Tumors in Male Rats  
Administered N-Nitrosodiphenylamine in the Diet (a)

(continued)

<u>Topography: Morphology</u>	<u>Matched Control</u>	<u>Low Dose</u>	<u>High Dose</u>
Pituitary: Chromophobe Adenoma (b)	9/18 (50)	9/47 (19)	7/47 (15)
P Values (c,d)	P = 0.020 (N)	P = 0.017 (N)	P = 0.006 (N)
Departure from Linear Trend (e)	P = 0.022		
Relative Risk (f)			
Lower Limit		0.383	0.298
Upper Limit		0.175	0.122
		0.930	0.772
Weeks to First Observed Tumor	100	100	100
Adrenal: Pheochromocytoma (b)	3/19 (16)	6/46 (13)	1/49 (2)
P Values (c,d)	P = 0.017 (N)	N.S.	N.S.
Relative Risk (f)			
Lower Limit		0.826	0.129
Upper Limit		0.204	0.003
		4.740	1.517
Weeks to First Observed Tumor	99	100	100

Table E1. Analyses of the Incidence of Primary Tumors in Male Rats  
Administered N-Nitrosodiphenylamine in the Diet (a)

(continued)		Matched Control		Low Dose		High Dose	
Topography:	Morphology						
Thyroid:	C-cell Adenoma (b)	3/19 (16)		1/49 (2)		2/48 (4)	
P Values (c,d)		N.S.		N.S.		N.S.	
Departure from Linear Trend (e)		P = 0.030 (N)					
Relative Risk (f)							
Lower Limit				0.129		0.264	
Upper Limit				0.003		0.024	
				1.517		2.160	
Weeks to First Observed Tumor		100		100		100	
Thyroid: Follicular-cell Carcinoma or Adenoma (b)		0/19 (0)		3/49 (6)		0/48 (0)	
P Values (c,d)		N.S.		N.S.		--	
Relative Risk (f)							
Lower Limit				Infinite		--	
Upper Limit				0.243		--	
				Infinite		--	
Weeks to First Observed Tumor		--		100		--	

Table El. Analyses of the Incidence of Primary Tumors in Male Rats  
Administered N-Nitrosodiphenylamine in the Diet (a)

(continued)		Matched Control	Low Dose	High Dose
Topography:	Morphology			
Pancreatic Islet:	Islet-cell			
Adenoma (b)		2/18 (11)	1/45 (2)	2/36 (6)
P Values (c,d)		N.S.	N.S.	N.S.
Relative Risk (f)				
Lower Limit			0.200	0.500
Upper Limit			0.004	0.040
			3.663	6.508
Weeks to First Observed Tumor		100	100	100
<hr/>				
Testis: Interstitial-cell				
Tumor (b)		15/19 (79)	38/50 (76)	38/49 (78)
P Values (c,d)		N.S.	N.S.	N.S.
Relative Risk (f)				
Lower Limit			0.963	0.982
Upper Limit			0.756	0.772
			1.401	1.419
Weeks to First Observed Tumor		88	95	100

Table El. Analyses of the Incidence of Primary Tumors in Male Rats  
Administered N-Nitrosodiphenylamine in the Diet (a)

(continued)

- (a) Dosed groups received 1,000 or 4,000 ppm.
- (b) Number of tumor-bearing animals/number of animals examined at site (percent).
- (c) Beneath the incidence of tumors in the control group is the probability level for the Cochran-Armitage test when P is less than 0.05; otherwise, not significant (N.S.) is indicated.  
Beneath the incidence of tumors in a dosed group is the probability level for the Fisher exact test for the comparison of that dosed group with the matched-control group when P is less than 0.05; otherwise, not significant (N.S.) is indicated.
- (d) A negative trend (N) indicates a lower incidence in a dosed group than in a control group.
- (e) The probability level for departure from linear trend is given when P is less than 0.05 for any comparison.
- (f) The 95 percent confidence interval of the relative risk between each dosed group and the control group.

Table E2. Analyses of the Incidence of Primary Tumors in Female Rats Administered N-Nitrosodiphenylamine in the Diet (a)

<u>Topography: Morphology</u>	<u>Matched Control</u>	<u>Low Dose</u>	<u>High Dose</u>
Integumentary System: Basal-cell Tumor of the Subcutaneous Tissue (b)	1/20 (5)	1/50 (2)	3/49 (6)
P Values (c,d)	N.S.	N.S.	N.S.
Relative Risk (f)			
Lower Limit		0.400	1.224
Upper Limit		0.005	0.108
		30.802	62.958
Weeks to First Observed Tumor	100	86	76
Integumentary System: Fibroma of the Subcutaneous Tissue (b)	0/20 (0)	1/50 (2)	3/49 (6)
P Values (c,d)	N.S.	N.S.	N.S.
Relative Risk (f)			
Lower Limit		Infinite	Infinite
Upper Limit		0.022	0.255
		Infinite	Infinite
Weeks to First Observed Tumor	--	100	97



Table E2. Analyses of the Incidence of Primary Tumors in Female Rats  
Administered N-Nitrosodiphenylamine in the Diet (a)

(continued)		Matched Control		Low Dose		High Dose	
Topography: Morphology							
Hematopoietic System:							
Granulocytic Leukemia (b)		3/20 (15)		0/50 (0)		0/49 (0)	
P Values (c,d)		P = 0.022 (N)		P = 0.021 (N)		P = 0.022 (N)	
Departure from Linear Trend (e)		P = 0.002					
Relative Risk (f)							
Lower Limit				0.000		0.000	
Upper Limit				0.000		0.000	
				0.659		0.673	
Weeks to First Observed Tumor		82		--		--	
Urinary Bladder:							
Transitional-cell Carcinoma (b)		0/18 (0)		0/48 (0)		40/49 (82)	
P Values (c,d)		P less than 0.001		--		P less than 0.001	
Departure from Linear Trend (e)		P = 0.042					
Relative Risk (f)							
Lower Limit				--		Infinite	
Upper Limit				--		5.314	
				--		Infinite	
Weeks to First Observed Tumor		--		--		69	

Table E2. Analyses of the Incidence of Primary Tumors in Female Rats  
Administered N-Nitrosodiphenylamine in the Diet (a)

(continued)				
<u>Topography: Morphology</u>	<u>Matched Control</u>	<u>Low Dose</u>	<u>High Dose</u>	
Pituitary: Chromophobe Adenoma (b)	8/19 (42)	13/49 (27)	12/48 (25)	
P Values (c,d)	N.S.	N.S.	N.S.	
Relative Risk (f)				
Lower Limit		0.630	0.594	
Upper Limit		0.306	0.281	
		1.512	1.445	
Weeks to First Observed Tumor	98	100	88	
Thyroid: C-cell Carcinoma or Adenoma (b)	1/20 (5)	5/49 (10)	3/48 (6)	
P Values (c,d)	N.S.	N.S.	N.S.	
Relative Risk (f)				
Lower Limit		2.041	1.250	
Upper Limit		0.254	0.110	
		94.440	64.251	
Weeks to First Observed Tumor	100	100	88	

Table E2. Analyses of the Incidence of Primary Tumors in Female Rats Administered N-Nitrosodiphenylamine in the Diet (a)

(continued)			
Topography: Morphology	Matched Control	Low Dose	High Dose
Mammary Gland: Fibroadenoma (b)	4/20 (20)	6/50 (12)	6/49 (12)
P Values (c,d)	N.S.	N.S.	N.S.
Relative Risk			
Lower Limit		0.600	0.612
Upper Limit		0.164	0.168
		2.659	2.710
Weeks to First Observed Tumor	82	88	100

(a) Dosed groups received 1,000 or 4,000 ppm.

(b) Number of tumor-bearing animals/number of animals examined at site (percent).

(c) Beneath the incidence of tumors in the control group is the probability level for the Cochran-Armitage test when P is less than 0.05; otherwise, not significant (N.S.) is indicated.

Beneath the incidence of tumors in a dosed group is the probability level for the Fisher exact test for the comparison of that dosed group with the matched-control group when P is less than 0.05; otherwise, not significant (N.S.) is indicated.

(d) A negative trend (N) indicates a lower incidence in a dosed group than in a control group.

(e) The probability level for departure from linear trend is given when P is less than 0.05 for any comparison.

(f) The 95 percent confidence interval of the relative risk between each dosed group and the control group.

## APPENDIX F

ANALYSES OF THE INCIDENCE OF PRIMARY TUMORS IN MICE  
ADMINISTERED N-NITROSODIPHENYLAMINE IN THE DIET



Table Fl. Analyses of the Incidence of Primary Tumors in Male Mice  
Administered N-Nitrosodiphenylamine in the Diet (a)

<u>Topography: Morphology</u>	<u>Matched Control</u>	<u>Low Dose</u>	<u>High Dose</u>
Lung: Alveolar/Bronchiolar Adenoma (b)	4/20 (20)	9/48 (19)	7/48 (15)
P Values (c,d)	N.S.	N.S.	N.S.
Relative Risk (f)			
Lower Limit		0.938	0.729
Upper Limit		0.307	0.216
		3.804	3.112
Weeks to First Observed Tumor	101	101	87
<hr/>			
Hematopoietic System: Lymphoma or Leukemia (b)	1/20 (5)	2/49 (4)	3/48 (6)
P Values (c,d)	N.S.	N.S.	N.S.
Relative Risk (f)			
Lower Limit		0.816	1.250
Upper Limit		0.046	0.110
		47.195	64.251
Weeks to First Observed Tumor	91	93	101



Table Fl. Analyses of the Incidence of Primary Tumors in Male Mice  
Administered N-Nitrosodiphenylamine in the Diet (a)

(continued)

<u>Topography:</u>	<u>Morphology</u>	<u>Matched Control</u>	<u>Low Dose</u>	<u>High Dose</u>
All Sites:	Hemangioma (b)	1/20 (5)	3/49 (6)	3/48 (6)
P Values (c,d)		N.S.	N.S.	N.S.
Relative Risk (f)				
Lower Limit			1.224	1.250
Upper Limit			0.108	0.110
			62.958	64.251
Weeks to First Observed Tumor		101	87	98
All Sites:	Hemangiosarcoma (b)	0/20 (0)	4/49 (8)	2/48 (4)
P Values (c,d)		N.S.	N.S.	N.S.
Relative Risk (f)				
Lower Limit			Infinite	Infinite
Upper Limit			0.394	0.128
			Infinite	Infinite
Weeks to First Observed Tumor		--	101	101

Table F1. Analyses of the Incidence of Primary Tumors in Male Mice  
Administered N-Nitrosodiphenylamine in the Diet (a)

(continued)		Matched Control		Low Dose		High Dose	
Topography:	Morphology						
All Sites:	Hemangioma	1/20 (5)		7/49 (14)		5/48 (10)	
	or Hemangiosarcoma (b)						
P Values (c,d)		N.S.		N.S.		N.S.	
Relative Risk (f)							
	Lower Limit			2.857		2.083	
	Upper Limit			0.411		0.259	
				125.833		96.358	
Weeks to First Observed Tumor		101		87		98	
Liver: Hepatocellular Carcinoma or Adenoma (b)		6/20 (30)		12/49 (24)		7/48 (15)	
P Values (c,d)		N.S.		N.S.		N.S.	
Relative Risk (f)							
	Lower Limit			0.816		0.486	
	Upper Limit			0.343		0.167	
				2.350		1.567	
Weeks to First Observed Tumor		101		72		101	

Table F1. Analyses of the Incidence of Primary Tumors in Male Mice  
Administered N-Nitrosodiphenylamine in the Diet (a)

(continued)

- (a) Dosed groups received 10,000 or 20,000 ppm.
- (b) Number of tumor-bearing animals/number of animals examined at site (percent).
- (c) Beneath the incidence of tumors in the control group is the probability level for the Cochran-Armitage test when P is less than 0.05; otherwise, not significant (N.S.) is indicated. Beneath the incidence of tumors in a dosed group is the probability level for the Fisher exact test for the comparison of that dosed group with the matched-control group when P is less than 0.05; otherwise, not significant (N.S.) is indicated.
- (d) A negative trend (N) indicates a lower incidence in a dosed group than in a control group.
- (e) The probability level for departure from linear trend is given when P is less than 0.05 for any comparison.
- (f) The 95 percent confidence interval of the relative risk between each dosed group and the control group.

Table F2. Analyses of the Incidence of Primary Tumors in Female Mice Administered N-Nitrosodiphenylamine in the Diet (a)

<u>Topography: Morphology</u>	<u>Matched Control</u>	<u>Low Dose</u>	<u>High Dose</u>
Lung: Alveolar/Bronchiolar Carcinoma or Adenoma (b)	3/20 (15)	11/47 (23)	5/38 (13)
P Values (c,d)	N.S.	N.S.	N.S.
Relative Risk (f)			
Lower Limit		1.560	0.877
Upper Limit		0.480	0.195
		8.051	5.213
Weeks to First Observed Tumor	101	80	101
Hematopoietic System: Lymphoma or Leukemia (b)	1/20 (5)	4/50 (8)	4/41 (10)
P Values (c,d)	N.S.	N.S.	N.S.
Relative Risk (f)			
Lower Limit		1.600	1.951
Upper Limit		0.175	0.214
		77.169	93.623
Weeks to First Observed Tumor	77	95	63

Table F2. Analyses of the Incidence of Primary Tumors in Female Mice Administered N-Nitrosodiphenylamine in the Diet (a)

(continued)

<u>Topography: Morphology</u>	<u>Matched Control</u>	<u>Low Dose</u>	<u>High Dose</u>
All Sites: Hemangioma (b)	0/20 (0)	1/50 (2)	3/41 (7)
P Values (c,d)	N.S.	N.S.	N.S.
Relative Risk (f)			
Lower Limit		Infinite	Infinite
Upper Limit		0.022	0.305
		Infinite	Infinite
Weeks to First Observed Tumor	--	101	94
Liver: Hepatocellular Carcinoma or Adenoma (b)	3/20 (15)	7/49 (14)	4/38 (11)
P Values (c,d)	N.S.	N.S.	N.S.
Relative Risk (f)			
Lower Limit		0.952	0.702
Upper Limit		0.250	0.134
		5.317	4.432
Weeks to First Observed Tumor	101	100	101

Table F2. Analyses of the Incidence of Primary Tumors in Female Mice Administered N-Nitrosodiphenylamine in the Diet (a)

(continued)

- 
- (a) Dosed groups received time-weighted average doses of 2,315 or 5,741 ppm.
  - (b) Number of tumor-bearing animals/number of animals examined at site (percent).
  - (c) Beneath the incidence of tumors in the control group is the probability level for the Cochran-Armitage test when P is less than 0.05; otherwise, not significant (N.S.) is indicated. Beneath the incidence of tumors in a dosed group is the probability level for the Fisher exact test for the comparison of that dosed group with the matched-control group when P is less than 0.05; otherwise, not significant (N.S.) is indicated.
  - (d) A negative trend (N) indicates a lower incidence in a dosed group than in a control group.
  - (e) The probability level for departure from linear trend is given when P is less than 0.05 for any comparison.
  - (f) The 95 percent confidence interval of the relative risk between each dosed group and the control group.





Review of the Bioassay of N-Nitrosodiphenylamine\* for Carcinogenicity  
by the Data Evaluation/Risk Assessment Subgroup  
of the Clearinghouse on Environmental Carcinogens

December 13, 1978

The Clearinghouse on Environmental Carcinogens was established in May, 1976, in compliance with DHEW Committee Regulations and the Provisions of the Federal Advisory Committee Act. The purpose of the Clearinghouse is to advise the Director of the National Cancer Institute on the Institute's bioassay program to identify and evaluate chemical carcinogens in the environment to which humans may be exposed. The members of the Clearinghouse have been drawn from academia, industry, organized labor, public interest groups, and State health officials. Members have been selected on the basis of their experience in carcinogenesis or related fields and, collectively, provide expertise in chemistry, biochemistry, biostatistics, toxicology, pathology, and epidemiology. Representatives of various Governmental agencies participate as ad hoc members. The Data Evaluation/Risk Assessment Subgroup of the Clearinghouse is charged with the responsibility of providing a peer review of reports prepared on NCI-sponsored bioassays of chemicals studied for carcinogenicity. It is in this context that the below critique is given on the bioassay of N-Nitrosodiphenylamine.

The primary reviewer for the report on the bioassay of N-Nitrosodiphenylamine agreed with the conclusion that the compound was carcinogenic in treated rats but was not in treated mice, under the conditions of test. After a brief description of the experimental design, he said that there were no outstanding shortcomings worth noting. Based on the findings, he said that N-Nitrosodiphenylamine may be a potential human carcinogen.

The secondary reviewer raised a question as to whether the bladder tumors in the treated rats were related to the presence of calculi. NCI pathologists responded that no calculi were reported by the examining pathologist and, at this point, there was no way of determining if they were specifically looked for. Since a carcinogenic effect was demonstrated, one Subgroup member commented that the report should stand on its own even though the mechanism by which the bladder tumors were induced is unknown.

A Subgroup member said that N-Nitrosodiphenylamine is a classical, non-biologically active nitrosamine. He suggested that the test compound may have nitrosated an amine present in the food which resulted in the formation of a carcinogenic nitrosamine whose target organ was the bladder. Because of the possibility, this Subgroup member urged

great caution in the interpretation of the results of the study for man. He recommended that the compound be retested using a diet free of nitrosatable amines.

After discussion regarding the framing of an appropriate motion, it was moved that the report on the bioassay of N-Nitrosodiphenylamine be accepted with the addition of comments to the Report's Summary section concerning: 1) the unknown role of calculi in the etiology of the bladder cancer in treated rats, because of the lack of knowledge as to whether they were present and; 2) the uncertainty as to whether the test compound reacted with a nitrosatable amine(s) in the diet to form a carcinogenic nitrosamine responsible for the induction of the bladder tumors. The motion was seconded and approved unanimously.


Clearinghouse Members Present:

Arnold L. Brown (Chairman), University of Wisconsin Medical School  
Joseph Highland, Environmental Defense Fund  
William Lijinsky, Frederick Cancer Research Center  
Henry Pitot, University of Wisconsin Medical Center  
Verne A. Ray, Pfizer Medical Research Laboratory  
Verald K. Rowe, Dow Chemical USA  
Michael Shimkin, University of California at San Diego  
Louise Strong, University of Texas Health Sciences Center  
Kenneth Wilcox, Michigan State Health Department

- 
- \* Subsequent to this review, changes may have been made in the bioassay report either as a result of the review or other reasons. Thus, certain comments and criticisms reflected in the review may no longer be appropriate.









GAYLORD			PRINTED IN U.S.A.

GAYLORD

PRINTED IN U. S. A.



4 0144 2792

NIH Library, Building 10  
National Institutes of Health  
Bethesda, Md. 20205



<http://nihlibrary.nih.gov>

---

10 Center Drive  
Bethesda, MD 20892-1150  
301-496-1080

NIH LIBRARY



3 1496 00123 5376

~~DEC 27 1988~~

DHEW Publication No. (NIH) 79-1720